arXiv:1209.3922v1 [math.AG] 18 Sep 2012

TORIC SHEAVES ON WEIGHTED PROJECTIVE PLANES

AMIN GHOLAMPOUR, YUNFENG JIANG AND MARTIJN KOOL

Abstract. We give an explicit description of toric sheaves on the weighted projective plane $\mathbb{P}(a,b,c)$ viewed as a toric Deligne-Mumford stack. The stack $\mathbb{P}(a,b,c)$ has an action of the torus $\mathbf{T} = \mathbb{C}^{*2}$ and by a toric sheaf we mean a coherent sheaf together with a lift of the torus action. The integers (a,b,c) are not necessarily chosen coprime or mutually coprime allowing for gerbe and root stack structures.

This is a first step in an attempt to give an explicit description of toric sheaves on any smooth toric Deligne-Mumford stack. Specializing the description to torsion free sheaves gives a combinatorial picture with various new features, which do not appear in the case of smooth toric varieties.

As an application, we describe the fixed point locus of the moduli scheme of stable rank 1 and 2 torsion free sheaves on $\mathbb{P}(a,b,c)$ with fixed K-group class. Summing over all K-group classes, we obtain formulae for generating functions of the topological Euler characteristics of these moduli schemes.

1. Introduction

A toric sheaf on a smooth toric variety X is a coherent sheaf together with a lift of the action of the torus $\mathbf{T}$ on X. More precisely, a toric sheaf consists of data $(\mathcal{F}, \Phi)$ where $\mathcal{F}$ is a coherent sheaf on X and Φ is a $\mathbf{T}$-equivariant structure on $\mathcal{F}$. Toric sheaves on smooth toric varieties have been well-studied, e.g. [Kly1, Per1] among others. The idea is to cover X by affine open $\mathbf{T}$-invariant subsets $U_\sigma \cong \mathbb{C}^d$, corresponding to the cones σ of maximal dimension in the fan Σ of X and restrict $(\mathcal{F}, \Phi)$ to each of these open subsets. Over these open subsets, the toric sheaf is given by a module with an $X(\mathbf{T})$-grading, where $X(\mathbf{T})$ is the character group of $\mathbf{T}$. This local data becomes particularly explicit in the case $\mathcal{F}$ is in addition locally free, reflexive, torsion free or just pure. Supplementing this local data with gluing conditions gives an explicit combinatorial description of $(\mathcal{F}, \Phi)$ in terms of the fan Σ of X. This can be used to compute Chern classes of $(\mathcal{F}, \Phi)$ and describe moduli of toric sheaves [Kly2, Per2, Pay, Koo1]. This in turn can be used to compute the Euler characteristic of the whole moduli space of stable sheaves on X using localization [Kly2, Koo2].

Our goal is to generalize the above to toric Deligne-Mumford stacks[1] $\mathfrak{X}$ introduced by [BCS]. Such stacks are described by a stacky fan $\mathbf{\Sigma}$ [BCS, FMN]. Again, one can consider the cones σ of maximal dimension in $\mathbf{\Sigma}$, which gives a cover by open $\mathbf{T}$-invariant substacks $\mathfrak{U}_\sigma \cong [\mathbb{C}^d/N(\sigma)]$ [BCS, Prop. 4.1]. Here $N(\sigma)$ is a finite abelian group acting on $\mathbb{C}^d$. We can then restrict a toric sheaf $(\mathcal{F}, \Phi)$ on $\mathfrak{X}$ to each of these open substacks. Viewing these open substacks as groupoids, we get similar data as before, but with two new phenomena:

i) The action of the torus $\mathbf{T}$ on $\mathbb{C}^d$ can be *non-primitive*: on a smooth toric variety, one can always choose coordinates on U_σ such that the action becomes $(\lambda_1, \ldots, \lambda_d) \cdot (x_1, \ldots, x_d) = (\lambda_1 x_1, \ldots, \lambda_d x_d)$, but in the current situation higher powers of λ_i can occur.
ii) The sheaf $\mathcal{F}|_{\mathfrak{U}_\sigma}$ is described by a module carrying a grading by $X(\mathbf{T})$ and each weight space carries a further *fine grading* by the character group $X(N(\sigma))$ of $N(\sigma)$.

This local data becomes particularly explicit when $\mathcal{F}$ is in addition locally free, reflexive, torsion free etc.

In order to recover $(\mathcal{F}, \Phi)$, one has to impose gluing conditions on the local data. Rather than trying to work this out for a general toric DM stack, we consider the specific example of the stacky weighted projective plane $\mathbb{P} := \mathbb{P}(a, b, c)$ for *any* positive integers a, b, c. In the case of smooth toric varieties, the gluing proceeds by intersecting any two open affine invariant subsets U_σ, U_τ and matching the local data. This involves pulling back the local data along *open immersions*. This times, we have to "intersect" any two open substacks $\mathfrak{U}_\sigma \cong [\mathbb{C}^d/N(\sigma)]$, $\mathfrak{U}_\tau \cong [\mathbb{C}^d/N(\tau)]$ and pull-back the local data to the stack theoretic intersection. At the level of groupoids, this turns out to correspond to pulling back the local data along certain *étale morphisms*. Working this out for $\mathbb{P}$, we get an explicit description of toric sheaves on stacky weighted projective planes. This can be used to study toric vector bundles and toric torsion free sheaves on $\mathbb{P}$ and describe their moduli.

As an application, we first describe the equivariant Picard group $\mathrm{Pic}^{\mathbf{T}}(\mathbb{P})$. Next, we consider the moduli spaces of stable rank 1 and 2 torsion free sheaves on $\mathbb{P}$. These moduli spaces[2], which are in fact schemes, have been introduced by [Nir] for any rank and any (not necessarily toric) projective DM stack $\mathbb{P}$. We briefly recall the relevant parts of Nironi's construction at the beginning of

---

[1] We only consider the case in which the stack $\mathfrak{X}$ is smooth as in [BCS].

[2] As opposed to Nironi, we consider μ-stability instead of Gieseker stability and we fix the K-class of the sheaf instead of the modified Hilbert polynomial. Here μ is the linear term of the *modified* Hilbert polynomial divided by its quadratic term. This means μ depends on a choice of polarization and *generating sheaf*.

Section 7. The action of the torus $\mathbf{T}$ on $\mathbb{P}$ lifts to the moduli space and we use the explicit description of toric sheaves on $\mathbb{P}$ to compute the fixed point locus and its topological Euler characteristic. The rank 1 case leads to counting of colored partitions (e.g. as in [DS]). In the rank 2 case, we get interesting new generating functions. Nironi's moduli spaces a priori depend on a choice of polarization and *generating sheaf*. On $\mathbb{P}$, the dependence on the polarization is trivial. However, the generating sheaf does play an important role.

Convention. Throughout this paper we fix $\mathbb{P} := \mathbb{P}(a, b, c)$ with a, b, c *arbitrary* positive integers.

Acknowledgments. We thank Paul Johnson and Benjamin Young for useful discussions on counting of colored partitions. We also thank Richard Thomas for helpful discussions. The third author is supported by EPSRC grant EP/G06170X/1, "Applied derived categories".

2. G-EQUIVARIANT SHEAVES ON DM STACKS WITH G-ACTION

Let $\mathfrak{X}$ be a DM stack[3]. A (quasi-)coherent sheaf $\mathcal{F}$ on $\mathfrak{X}$ is given by specifying a (quasi-)coherent sheaf $\mathcal{F}_U$ over each atlas $U \to \mathfrak{X}$ together with compatibility isomorphisms between the $\mathcal{F}_U$ [Vis, Def. 7.18]. We are interested in the case $\mathfrak{X}$ carries the action of an affine algebraic group G and $\mathcal{F}$ has a G-equivariant structure. Note that the affine algebraic group G is taken to be a variety so is non-stacky.

Definition 2.1. [Rom, Def. 1.3, 4.3] Let $\mathfrak{X}$ be a DM stack and G an affine algebraic group with multiplication $\mu : G \times G \to G$, inverse map $\iota : G \to G$ and identity $e : \operatorname{Spec} \mathbb{C} \to G$. A *$G$-action* on $\mathfrak{X}$ is a morphism $\boldsymbol{\sigma} : G \times \mathfrak{X} \to \mathfrak{X}$ together with 2-isomorphisms $\boldsymbol{\alpha}$ and $\boldsymbol{a}$

$$\begin{tikzcd}
G \times G \times \mathfrak{X} \arrow[r, "\mu\times\mathbf{id}_{\mathfrak{X}}"] \arrow[d, "\mathrm{id}_G\times\boldsymbol{\sigma}"'] & G \times \mathfrak{X} \arrow[d, "\boldsymbol{\sigma}"] \\
G \times \mathfrak{X} \arrow[r, "\boldsymbol{\sigma}"'] \arrow[ur, Rightarrow, "\boldsymbol{\alpha}"] & \mathfrak{X}
\end{tikzcd}
\qquad
\begin{tikzcd}
\operatorname{Spec} \mathbb{C} \times \mathfrak{X} \arrow[r, "e\times\mathbf{id}_{\mathfrak{X}}"] \arrow[dr, equal] & G \times \mathfrak{X} \arrow[d, "\boldsymbol{\sigma}"] \arrow[dl, Rightarrow, "\boldsymbol{a}"] \\
 & \mathfrak{X},
\end{tikzcd}$$

where the 2-isomorphisms $\boldsymbol{\alpha}$, $\boldsymbol{a}$ satisfy some natural compatibility conditions (see [Rom, Def. 1.3]).

Next, let $\mathcal{F}$ be a quasi-coherent sheaf on $\mathfrak{X}$. Denote by $\boldsymbol{p}_2 : G \times \mathfrak{X} \to \mathfrak{X}$ and $\boldsymbol{p}_{23} : G \times G \times \mathfrak{X} \to G \times \mathfrak{X}$ projection on the second and last two factors.

---

[3] All our stacks and schemes are defined over $\mathbb{C}$. DM stacks are defined over the étale site.

A *G-equivariant structure* on $\mathcal{F}$ is an isomorphism $\Phi : \boldsymbol{\sigma}^*\mathcal{F} \to \boldsymbol{p}_2^*\mathcal{F}$ satisfying the cocycle identity

$$\boldsymbol{p}_{23}^*\Phi \circ (\mathrm{id}_G \times \boldsymbol{\sigma})^*\Phi = (\mu \times \mathbf{id}_{\mathfrak{X}})^*\Phi.$$

When Φ is a G-equivariant structure on $\mathcal{F}$, we refer to the pair $(\mathcal{F}, \Phi)$ as a *G-equivariant sheaf.* A *G-equivariant morphism* between G-equivariant sheaves $(\mathcal{F}, \Phi)$, $(\mathcal{G}, \Psi)$ is a morphism $\theta : \mathcal{F} \to \mathcal{G}$ satisfying $\boldsymbol{p}_2^*\theta \circ \Phi = \Psi \circ \boldsymbol{\sigma}^*\theta$. $\oslash$

We leave various straight-forward notions such as "G-equivariant morphisms between DM stacks with G-action", "G-invariant open substacks" and "pull-back of G-equivariant sheaves along G-equivariant morphisms" to the reader.

Suppose $\{\mathfrak{U}_i\}$ is a cover by G-invariant open substacks of a DM stack $\mathfrak{X}$ with G-action. We want to think of G-equivariant quasi-coherent sheaves on $\mathfrak{X}$ as G-equivariant quasi-coherent sheaves on the $\mathfrak{U}_i$ together with gluing conditions. The idea is that in applications, G-equivariant quasi-coherent sheaves on the $\mathfrak{U}_i$ could be much easier to describe, e.g. when the $\mathfrak{U}_i$ are quotient stacks. We use the notation $|_i$, $|_{ij}$, $|_{ijk}$ for restriction $|_{\mathfrak{U}_i}$, $|_{\mathfrak{U}_{ij}}$, $|_{\mathfrak{U}_{ijk}}$, where $\mathfrak{U}_{ij} := \mathfrak{U}_i \times_{\mathfrak{X}} \mathfrak{U}_j$ and $\mathfrak{U}_{ijk} := \mathfrak{U}_i \times_{\mathfrak{X}} \mathfrak{U}_j \times_{\mathfrak{X}} \mathfrak{U}_k$. Using the definition of a quasi-coherent sheaf on a DM stack [Vis, Def. 7.18] and [Har, Exc. II.1.22] on atlases, gives:

Proposition 2.2. *Let $\mathfrak{X}$ be a DM stack with G-action and let $\{\mathfrak{U}_i\}$ be a cover by G-invariant open substacks. Let $\{(\mathcal{F}_i, \Phi_i)\}$ be a collection of G-equivariant quasi-coherent sheaves on the $\mathfrak{U}_i$ and assume there are G-equivariant isomorphisms $\phi_{ij} : \mathcal{F}_i|_{ij} \to \mathcal{F}_j|_{ij}$ satisfying $\phi_{ii} = \mathrm{id}$ and the cocycle identity $\phi_{ik}|_{ijk} = \phi_{jk}|_{ijk} \circ \phi_{ij}|_{ijk}$. Then there exists a unique G-equivariant quasi-coherent sheaf $(\mathcal{F}, \Phi)$ on $\mathfrak{X}$ together with G-equivariant isomorphisms $\phi_i : \mathcal{F}|_i \to \mathcal{F}_i$ satisfying $\phi_j|_{ij} = \phi_{ij} \circ \phi_i|_{ij}$.*

Let H be an affine algebraic group acting on a scheme X and assume the stabilizers of the $\mathbb{C}$-points are reduced and finite. Then $\mathfrak{X} = [X/H]$ is a DM stack [Vis, Ex. 7.17] and we refer to such stacks as quotient DM stacks. For later use, it is more convenient to work with the natural groupoid presentation of a quotient stack $[X/H]$

$$\begin{tikzcd} H \times X \arrow[r, shift left, "p_2"] \arrow[r, shift right, "\tau"'] & X, \end{tikzcd}$$

where p_2 (the source map of the groupoid) is projection to the first factor and τ (the target map of the groupoid) is the given action of H on X [Vis, Sect. 7]. We suppress notation of the other three (obvious) structure morphisms.

Proposition 2.3. [Vis, Ex. 7.21] *The category of quasi-coherent sheaves on a quotient DM stack $[X/H]$ is equivalent to the category of H-equivariant quasi-coherent sheaves on X.*

We want to generalise this to the G-equivariant setting. We start with the following lemma.

Lemma 2.4. *Let $\mathfrak{X} = [X/H]$ be a quotient DM stacks. Any G-action on X commuting with H induces a canonical G-action on $\mathfrak{X}$.*

Proof. Let $\sigma : G \times X \to X$ be the G-action on X. The stack $G \times \mathfrak{X}$ admits a groupoid presentation

$$\begin{tikzcd} G \times H \times X \arrow[r, shift left, "\mathrm{id}_G \times p_2"] \arrow[r, shift right, "\mathrm{id}_G \times \tau"'] & G \times X, \end{tikzcd}$$

which is induced from the natural groupoid presentation of $\mathfrak{X}$. The other structure maps are the obvious maps. Let $\sigma_1 := p_2 \times (\sigma \circ p_{13}) : G \times H \times X \to H \times X$ and $\sigma_0 := \sigma$. The commutativity of the actions of H and G on X implies that (σ_1, σ_0) is a morphism of groupoids

$$\begin{tikzcd} G \times H \times X \arrow[r, shift left, "\mathrm{id}_G \times p_2"] \arrow[r, shift right, "\mathrm{id}_G \times \tau"'] \arrow[d, "\sigma_1"'] & G \times X \arrow[d, "\sigma_0"] \\ H \times X \arrow[r, shift left, "p_2"] \arrow[r, shift right, "\tau"'] & X, \end{tikzcd}$$

which gives rise to the morphism between the associated stacks. It is straightforward to check that this morphism indeed defines a G-action on $\mathfrak{X}$ as in Definition 2.1 with $\boldsymbol{\alpha}$ and $\boldsymbol{a}$ both taken to be the identity. □

Given a scheme X with commuting G- and H-actions and a quasi-coherent sheaf $\mathcal{F}$ on X, we can consider commuting G- and H-equivariant structures Φ and Ψ on $\mathcal{F}$. Denoting the actions by $\sigma : G \times X \to X$, $\tau : H \times X \to X$ and projections by $p_{13} : G \times H \times X \to G \times X$ and $p_{23} : G \times H \times X \to H \times X$, this means

$$p_{13}^*\Phi \circ (\mathrm{id}_H \times \sigma)^*\Psi = p_{23}^*\Psi \circ (\mathrm{id}_G \times \tau)^*\Phi. \tag{1}$$

Triples $(\mathcal{F}, \Phi, \Psi)$ with Φ and Ψ commuting can be made into a category by considering morphisms which intertwine both the G- and H-equivariant structure.

Proposition 2.5. *Let $\mathfrak{X} = [X/H]$ be a quotient DM stack with G-action coming from a G-action on X commuting with H. Then the category of G-equivariant quasi-coherent sheaves on $\mathfrak{X}$ is equivalent to the category of quasi-coherent sheaves on X with commuting G- and H-equivariant structures.*

Proof. The proof is a straightforward application of the equivalence of categories in Proposition 2.3. We denote the G-action on X by σ and the H-action on X by τ and use p_i, p_{ij} for the various projections as before.

Let $\boldsymbol{\sigma}, \boldsymbol{p} : G \times \mathfrak{X} \to \mathfrak{X}$ be the G-action and projection. These two morphisms of stacks are obtained from the corresponding morphisms $(\sigma_1, \sigma_0) = (p_2 \times (\sigma \circ p_{13}), \sigma)$ and $(p_1, p_0) = (p_{23}, p_2)$ of the groupoid presentations

$$\begin{tikzcd}
G \times H \times X \arrow[r, shift left, "s'"] \arrow[r, shift right, "t'"'] \arrow[d, shift right, "\sigma_1"'] \arrow[d, shift left, "p_1"] & G \times X \arrow[d, shift right, "\sigma_0"'] \arrow[d, shift left, "p_0"] \\
H \times X \arrow[r, shift left, "s"] \arrow[r, shift right, "t"'] & X,
\end{tikzcd}$$

where $s = p_2$, $t = \tau$, $s' = \mathrm{id}_G \times p_2$ and $t' = \mathrm{id}_G \times \tau$. By Proposition 2.3, giving a G-equivariant quasi-coherent sheaf $\mathcal{F}$ on $\mathfrak{X}$ is equivalent to giving a quasi-coherent sheaf $\mathcal{F}_X$ on X together with four isomorphisms

$$\Psi : t^*\mathcal{F}_X \to s^*\mathcal{F}_X, \quad A : t'^*\sigma_0^*\mathcal{F}_X \to s'^*\sigma_0^*\mathcal{F}_X,$$
$$B : t'^*p_0^*\mathcal{F}_X \to s'^*p_0^*\mathcal{F}_X, \quad \Phi : \sigma_0^*\mathcal{F}_X \to p_0^*\mathcal{F}_X,$$

such that:

i) Ψ satisfies the cocycle identity,
ii) $A = \sigma_1^*\Psi$, $B = p_1^*\Psi$,
iii) $s'^*\Phi \circ A = B \circ t'^*\Phi$,
iv) Φ satisfies the cocycle identity.

The isomorphism Ψ (with cocycle condition) defines an H-equivariant structure on $\mathcal{F}_X$. The isomorphisms A, B together with condition ii) and the sheaves $\sigma_0^*\mathcal{F}_X$ and $p_0^*\mathcal{F}_X$ correspond to the quasi-coherent sheaves $\boldsymbol{\sigma}^*\mathcal{F}$ and $\boldsymbol{p}^*\mathcal{F}$ on the stack $G \times \mathfrak{X}$. The isomorphism Φ with condition iii) corresponds to an isomorphism $\boldsymbol{\sigma}^*\mathcal{F} \cong \boldsymbol{p}^*\mathcal{F}$ on the stack $G \times \mathfrak{X}$. It satisfies the cocycle identity if and only if iv) holds. This defines a G-equivariant structure on $\mathcal{F}_X$. This finishes the proof by noting that the condition iii) above is equivalent to Equation (1). □

Consider the setting in the previous proposition. If in addition $X = \mathrm{Spec}\, A$ is affine, G and H are diagonalizable, then we get a further equivalence with the category of A-modules with $X(G) \times X(H)$-grading as follows. The global section functor $H^0(X, \cdot)$ gives an equivalence between the category of quasi-coherent sheaves on X and the category A-modules. If a quasi-coherent sheaf $\mathcal{F}$ has in addition a G- and H-equivariant structure, then this can be used to define a linear G and H-action on $H^0(X, \mathcal{F})$ as follows. Let $i_g : \mathrm{Spec}\, \mathbb{C} \hookrightarrow G$ be inclusion of an element g into G and set $\Phi_g := i_g^*\Phi$. Then for any $s \in H^0(X, \mathcal{F})$, we define $g \cdot s := \Phi_{g^{-1}}((g^{-1})^* s) \in H^0(X, \mathcal{F})$, where $(g^{-1})^* s \in H^0(X, (g^{-1})^* \mathcal{F})$ is the canonical lift (and similarly for H). Note that Φ and Ψ commute, so the G- and H-action on $H^0(X, \mathcal{F})$ commute. Since G is diagonalizable, the module $H^0(X, \mathcal{F})$ decomposes into G-eigenspaces

$$H^0(X, \mathcal{F}) = \bigoplus_{\chi \in X(G)} H^0(X, \mathcal{F})_\chi.$$

Each $H^0(X, \mathcal{F})_m$ has an induced H-action. Since H is diagonalizable, we get a further decomposition into eigenspaces

$$H^0(X, \mathcal{F}) = \bigoplus_{\chi \in X(G)} \bigoplus_{\psi \in X(H)} H^0(X, \mathcal{F})_{\chi, \psi}.$$

For H trivial, Kaneyama [Kan] proves $H^0(X, \cdot)$ gives an equivalence between the category of G-equivariant quasi-coherent sheaves on X and the category of $X(G)$-graded A-modules (see also [Per3, Prop. 2.31]). It is not hard to show that in our current setting, $H^0(X, \cdot)$ gives an equivalence between the category of quasi-coherent sheaves on X with commuting G- and H-equivariant structures and the category of $X(G) \times X(H)$-graded A-modules. We find it convenient in our applications to refer to the $X(H)$-grading as the *fine grading.* Summarizing:

Corollary 2.6. *Let $\mathfrak{X} = [(\mathrm{Spec}\ A)/H]$ be a quotient DM stack with G-action coming from a G-action on* Spec A *commuting with H. Assume G and H are diagonalizable. Then the category of quasi-coherent sheaves on $\mathfrak{X}$ with commuting G- and H-equivariant structures is equivalent to the category of A-modules with an $X(G)$-grading and $X(H)$-fine grading.*

The idea of this corollary is that in applications, moduli of finitely generated A-modules with $X(G)$-grading and $X(H)$-fine grading can be explicitly described.

3. Toric sheaves on affine toric DM stacks

Let $\mathbb{C}^d$ be affine space with linear $\mathbf{T} = \mathbb{C}^{*d}$-action. Let H be a finite abelian group. Suppose $\mathbb{C}^d$ also has an H-action and the $\mathbf{T}$ and H actions commute. In this section, we describe toric sheaves, i.e. $\mathbf{T}$-equivariant coherent sheaves, on quotient DM stacks $\mathfrak{X} = [\mathbb{C}^d/H]$. This boils down to applying Corollary 2.6 and repackaging the grading data somewhat. We also discuss what torsion free and reflexive sheaves look like in this picture.

Throughout this paper, we make the following notational conventions. Let D be any diagonalizable group. Then $D \cong \mathbb{C}^{*d} \times \boldsymbol{\mu}_{a_1} \times \cdots \times \boldsymbol{\mu}_{a_s}$, where $\boldsymbol{\mu}_{a_i}$ is the group of a_i-th roots of unity and the a_i's are prime powers. The numbers

d, a_i are unique with this property. The group operation of the latter group (and its factors) is written multiplicatively and its elements are denoted by $\lambda = (\lambda_1, \ldots, \lambda_d, \mu_1, \ldots, \mu_s)$. We denote the character group of D by $X(D)$. There is an isomorphism

$$X(D) \cong \mathbb{Z}^d \oplus \mathbb{Z}_{a_1} \oplus \cdots \oplus \mathbb{Z}_{a_s}.$$

The group operation of the group on the RHS (and its factors) is written additively and its elements are denoted by $m = (\ell_1, \ldots, \ell_d, l_1, \ldots, l_s)$. The isomorphism maps $m = (\ell_1, \ldots, \ell_d, l_1, \ldots, l_s)$ to the character

$$\chi(m) : D \longrightarrow \mathbb{C}^*, \ (\lambda_1, \ldots, \lambda_d, \mu_1, \ldots, \mu_s) \mapsto \lambda_1^{\ell_1} \cdots \lambda_d^{\ell_d} \mu_1^{l_1} \cdots \mu_s^{l_s}.$$

We often switch between additive notation $m, m', \ldots, +$ and multiplicative notation $\chi(m), \chi(m'), \ldots, \cdot$ for $X(D)$. With this notation $\chi(m)\chi(m') = \chi(m + m')$.

3.1. **Non-degenerate torus actions.**

We start with some elementary remarks on the action of $\mathbf{T}$ on $\mathbb{C}^d$.

Definition 3.1. Let $\mathbf{T}$ act linearly[4] on $\mathbb{C}^d$. Then there exist coordinates $x_1, \ldots, x_d$ and characters $\chi(m_1), \ldots, \chi(m_d)$ such that $\lambda \cdot x_i = \chi(m_i)(\lambda) x_i$ for all $i = 1, \ldots, d$ and $\lambda \in \mathbf{T}$. If the m_i's are dependent, then the action is said to be *degenerate*. If the action is non-degenerate and the m_i generate the lattice $X(\mathbf{T})$, then the action is said to be *primitive*. These notions do not depend on the choice of coordinates. $\oslash$

In the case of smooth toric varieties, the opens in the cover $\{U_\sigma\}$ corresponding to the cones σ of maximal dimension have primitive linear torus actions. In the case of smooth toric stacks, the opens in the cover $\{[\mathbb{C}^d/N(\sigma)]\}$ corresponding to cones σ of maximal dimension can have a torus action induced from a non-primitive linear torus action on $\mathbb{C}^d$. We are interested in non-degenerate linear torus actions.

Definition 3.2. Let $\mathbf{T}$ act linearly and non-degenerately on $\mathbb{C}^d$. Suppose the action is written as $\lambda \cdot x_i = \chi(m_i)(\lambda) x_i$. Note that this choice of coordinates is unique[5] up to scaling and reordering the x_i. The *box* associated to the action is the subset $\mathcal{B}_{\mathbf{T}} \subset X(\mathbf{T})$ of all elements of the form $\sum_i q_i m_i \in X(\mathbf{T})$ with $0 \leq q_1, \ldots, q_d < 1$ rational. Note that $\mathcal{B}_{\mathbf{T}} = 0$ if and only if the $\mathbf{T}$-action is primitive. We often denote an element $\sum_i q_i m_i \in \mathcal{B}_{\mathbf{T}}$ by $(q_1, \ldots, q_d) \in \mathbb{Q}^d$. $\oslash$

---

[4] It is conjectured that *any* $\mathbf{T}$-action on $\mathbb{C}^d$ is linear after a suitable change of coordinates (linearization conjecture). This has been proved for $d \leq 3$ [Gut, KR1, KR2].

[5] In the case of toric varieties, such a choice does not have to be made since $\mathbf{T} \subset \mathbb{C}^d$ and the action of $\mathbf{T}$ on $\mathbb{C}^d$ restricts to the canonical action $\mathbf{T} \times \mathbf{T} \to \mathbf{T}$.

For $\mathbb{C}^d$ with non-degenerate linear $\mathbf{T}$-action, the weight spaces $H^0(\mathbb{C}^d, \mathcal{O}_{\mathbb{C}^d})_m$ are 0 or 1-dimensional. The collection of $m \in X(\mathbf{T})$ for which $H^0(\mathbb{C}^d, \mathcal{O}_{\mathbb{C}^d})_m \neq 0$ forms a semigroup S. Writing the action as $\lambda \cdot x_i = \chi(m_i)(\lambda)x_i$, this is also the cone spanned by $m_1, \ldots, m_d \in X(\mathbf{T})$. The category of $\mathbf{T}$-equivariant quasi-coherent sheaves on $\mathbb{C}^d$ is equivalent to the category of $X(\mathbf{T})$-graded modules (Corollary 2.6). We repackage this data neatly in the same way Perling does in the case of toric varieties [Per1].

Suppose the action is written as $\lambda \cdot x_i = \chi(m_i)(\lambda)x_i$. Given a $\mathbf{T}$-equivariant quasi-coherent sheaf $\mathcal{F}$ on $\mathbb{C}^d$, let $H^0(\mathbb{C}^d, \mathcal{F}) = \bigoplus_{m \in X(\mathbf{T})} F(m)$ be the corresponding $X(\mathbf{T})$-graded module. This gives us a collection of vector spaces $\{F(m)\}_{m \in X(\mathbf{T})}$. Multiplication by x_i gives linear maps

$$\chi_i(m) : F(m) \longrightarrow F(m + m_i),$$

for all $m \in X(\mathbf{T})$. These maps satisfy

$$\chi_j(m + m_i) \circ \chi_i(m) = \chi_i(m + m_j) \circ \chi_j(m), \tag{2}$$

for any $m \in X(\mathbf{T})$ and $i, j = 1, \ldots, d$. Abstractly, we refer to a collection of vector spaces $\{F(m)\}_{m \in X(\mathbf{T})}$ and linear maps $\{\chi_i(m)\}_{m \in X(\mathbf{T}), i=1,\ldots,d}$ satisfying (2) as an *S-family* $\hat{F}$:

Definition 3.3. Let $X(\mathbf{T})$ be the character group of an algebraic torus $\mathbf{T}$ of dimension d. Suppose we are given $\mathbb{Z}$-independent elements $m_1, \ldots, m_d \in X(\mathbf{T})$. An *$S$-family* $\hat{F}$ consists of the following data: a collection of vector spaces $\{F(m)\}_{m \in X(\mathbf{T})}$ and linear maps $\{\chi_i(m)\}_{m \in X(\mathbf{T}), i=1,\ldots,d}$ satisfying (2). (The letter S stands for the semigroup given by the cone generated by $m_1, \ldots, m_d \in X(\mathbf{T})$.) A *morphism of S-families* $\hat{F}$, $\hat{G}$ is a family $\hat{\phi}$ of linear maps $\{\phi(m) : F(m) \to G(m)\}_{m \in X(\mathbf{T})}$ commuting with the $\chi_i(m)$'s. ⊘

The S-family $\hat{F}$ associated to a $\mathbf{T}$-equivariant quasi-coherent sheaf $\mathcal{F}$ contains all the data of the graded module $H^0(\mathcal{F})$. It is easy to see that the category of $\mathbf{T}$-equivariant quasi-coherent sheaves on $\mathbb{C}^d$ is equivalent to the category of S-families. The following is also obvious.

Proposition 3.4. *Let $\mathbf{T}$ act non-degenerately on $\mathbb{C}^d$. Then each $\mathbf{T}$-equivariant quasi-coherent sheaf $\mathcal{F}$ on $\mathbb{C}^d$ with corresponding S-family $\hat{F}$ decomposes equivariantly according to the box elements*

$$\mathcal{F} \cong \bigoplus_{b \in \mathcal{B}_{\mathbf{T}}} {}_b\mathcal{F}, \ \ \hat{F} \cong \bigoplus_{b \in \mathcal{B}_{\mathbf{T}}} {}_b\hat{F}.$$

Proof. It suffices to show the statement for S-families. Take an element $b \in \mathcal{B}_{\mathbf{T}}$ then define ${}_b\hat{F}$ to be the S-subfamily consisting of all vector spaces $F(m + b)$ where $m = \sum_i \ell_i m_i$ for any $\ell_i \in \mathbb{Z}$. □

Summary. A $\mathbf{T}$-equivariant quasi-coherent sheaf on affine space $\mathbb{C}^d$ with non-degenerate linear $\mathbf{T}$-action is just a family of vector spaces indexed by the lattice points of $X(\mathbf{T})$ and (compatible linear) maps between them encoding the module structure. When the action is non-primitive, the sheaf decomposes according to the box elements $b \in \mathcal{B}_{\mathbf{T}}$.

We end this section by mentioning what the S-family of a coherent, torsion free and reflexive sheaf looks like. Let $\mathcal{F}$ be a $\mathbf{T}$-equivariant quasi-coherent sheaf on $\mathbb{C}^d$ with S-family $\hat{F}$. Any $m \in X(\mathbf{T})$ can be uniquely written as $b + \sum_i \ell_i m_i$ where the ℓ_i are integers. We often write ${}_bF(\ell_1, \ldots, \ell_d) := {}_bF(m)$. Moreover, the element $b \in \mathcal{B}_{\mathbf{T}}$ can be uniquely written as $\sum_i q_i m_i$ for rational numbers $0 \leq q_i < 1$ and we also often write ${}_{(q_1,\ldots,q_d)}F(\ell_1, \ldots, \ell_d) := {}_bF(\ell_1, \ldots, \ell_d)$. Similarly, the maps $\chi_i(b+m)$ are also denoted by ${}_b\chi_i(\ell_1, \ldots, \ell_d)$, ${}_{(q_1,\ldots,q_d)}\chi_i(\ell_1, \ldots, \ell_d)$ or simply x_i if we suppress the domain.

The sheaf $\mathcal{F}$ is *coherent* if and only if the corresponding module is finitely generated. In terms of $\hat{F}$ this means the following [Per1, Def. 5.10, Prop. 5.11]:

i) all vector spaces ${}_bF(\vec{\ell})$ are finite dimensional,
ii) for each $b \in \mathcal{B}_{\mathbf{T}}$ and $\ell_1, \ldots, \ell_d$ sufficiently negative ${}_bF(\vec{\ell}) = 0$,
iii) for each $b \in \mathcal{B}_{\mathbf{T}}$, there are only finitely many $\vec{L} \in \mathbb{Z}^d$ for which

$$ {}_bF(\vec{L}) \neq \mathrm{span}_{\mathbb{C}}\{x_1^{L_1-\ell_1} \cdots x_d^{L_d-\ell_d} s \mid s \in {}_bF(\vec{\ell}) \text{ and } L_i - \ell_i \geq 0 \text{ not all zero}\}. $$

We refer to $\mathbf{T}$-equivariant coherent sheaves as *toric sheaves*.

The sheaf $\mathcal{F}$ is *torsion free* if and only if it is coherent and all maps x_i are injective. This is proved in [Per1, Prop. 5.13] (or [Koo1, Prop. 2.8], which also discusses *pure* sheaves of lower dimension). Up to an equivalence of categories, this means we can take all x_i's to be inclusion. Decomposing $\hat{F} \cong \bigoplus_{b \in \mathcal{B}_{\mathbf{T}}} {}_b\hat{F}$, we conclude that a toric torsion free sheaf on $\mathbb{C}^d$ is specified by the following data:

i) a finite-dimensional vector spaces ${}_bF(\vec{\infty})$ for all $b \in \mathcal{B}_{\mathbf{T}}$,
ii) for each $b \in \mathcal{B}_{\mathbf{T}}$: a multifiltration $\{{}_bF(\vec{\ell})\}_{\vec{\ell} \in \mathbb{Z}^d}$ of ${}_bF(\vec{\infty})$ such that ${}_bF(\vec{\ell}) = {}_bF(\vec{\infty})$ for sufficiently large $\vec{\ell} \in \mathbb{Z}^d$.

The rank of such a torsion free sheaf is $\sum_{b \in \mathcal{B}_{\mathbf{T}}} \dim_{\mathbb{C}}({}_bF(\vec{\infty}))$.

How do we identify the *reflexive* sheaves among the torsion free sheaves? The following describe the S-families corresponding to toric reflexive sheaves. Suppose for each $b \in \mathcal{B}_{\mathbf{T}}$ we are given a finite-dimensional vector space ${}_bV(\infty)$ (not all zero) and a filtration

$$ \cdots \subset {}_bV(\ell - 1) \subset {}_bV(\ell) \subset {}_bV(\ell + 1) \subset \cdots , $$

such that ${}_bV(\ell) = 0$ for ℓ sufficiently small and ${}_bV(\ell) = {}_bV(\infty)$ for ℓ sufficiently large. Set

$$ {}_bF(\ell_1, \dots, \ell_d) = {}_bV(\ell_1) \cap \cdots \cap {}_bV(\ell_d) \subset {}_bV(\infty). \tag{3}$$

Such special types of multi-filtrations exactly correspond to the S-families of toric reflexive sheaves [Per1, Thm. 5.19].

3.2. **Toric sheaves on** $[\mathbb{C}^d/H]$**.** We now describe toric sheaves on the DM stack $[\mathbb{C}^d/H]$, where $\mathbf{T} = \mathbb{C}^{*d}$ and H is a finite abelian group, both acting on $\mathbb{C}^d$. We assume $\mathbf{T}$ acts linearly and non-degenerately and the actions of $\mathbf{T}$ and H commute. By Corollary 2.6, the category of $\mathbf{T}$-equivariant sheaves on $[\mathbb{C}^d/H]$ is equivalent to the category of $H^0(\mathbb{C}^d, \mathcal{O}_{\mathbb{C}^d})$-modules with $X(\mathbf{T})$-grading and $X(H)$-fine-grading.

Next, we choose coordinates x_i on $\mathbb{C}^d$ such that the $\mathbf{T}$-action is given by $\lambda \cdot x_i = \chi(m_i)(\lambda)x_i$. As mentioned before, this choice of coordinates is unique up to scaling and reordering the x_i. A $\mathbf{T}$-equivariant quasi-coherent sheaf $\mathcal{F}$ on $[\mathbb{C}^d/H]$ gives rise to the $H^0(\mathbb{C}^d, \mathcal{O}_{\mathbb{C}^d})$-module $H^0(\mathbb{C}^d, \mathcal{F})$ with $X(\mathbf{T})$-grading and $X(H)$-fine-grading. In turn, this provides us with an S-family $\hat{F}$ as described in Section 3.1. For each $m \in X(\mathbf{T})$, the vector space $F(m)$ has an $X(H)$-grading

$$F(m) = \bigoplus_{n \in X(H)} F(m)_n.$$

Since the actions of $\mathbf{T}$ and H commute, H acts by $h \cdot x_i = \chi(n_i)(h)x_i$ for some unique $n_i \in X(H)$. Each multiplication map $\chi_i(m)$ maps $F(\vec{\lambda})_n$ to $F(m + m_i)_{n+n_i}$. In other words, we have the following data:

i) a collection of vector spaces $\{F(m)_n\}_{m \in X(\mathbf{T}), n \in X(H)}$,

ii) a collection of linear maps $\{\chi_i(m) : F(m) \to F(m + m_i)\}_{i=1,\dots,d, m \in X(\mathbf{T})}$ satisfying

$$\begin{aligned} &\chi_i(m) : F(m)_n \longrightarrow F(m + m_i)_{n+n_i}, \\ &\chi_j(m + m_i) \circ \chi_i(m) = \chi_i(m + m_j) \circ \chi_j(m), \end{aligned}$$

for all $i, j = 1, \dots, d$, $m \in X(\mathbf{T})$ and $n \in X(H)$.

We refer to this data $\hat{F}$ as a *stacky S-family*. There is an obvious notion of morphism between stacky S-families: a morphism of S-families respecting the fine-grading. This repackaging can be summarized in the following proposition.

Proposition 3.5. *Let $\mathbf{T} = \mathbb{C}^{*d}$ and H be a finite abelian group acting on $\mathbb{C}^d$. Assume $\mathbf{T}$ acts linearly and non-degenerately and the actions of $\mathbf{T}$ and H commute. Then the category of $\mathbf{T}$-equivariant quasi-coherent sheaves on $[\mathbb{C}^d/H]$ is equivalent to the category of stacky S-families.*

Note that a stacky S-family decomposes according to the elements of the box $\mathcal{B}_{\mathbf{T}}$ as in Section 3.1. Coherent, torsion free and reflexive $\mathbf{T}$-equivariant quasi-coherent sheaves on $[\mathbb{C}^d/H]$ correspond to stacky S-families with underlying S-family satisfying the properties described in Section 3.1. We give two examples.

Example 1. Let $\mathbf{T} = \mathbb{C}^{*2}$ act by $(\kappa, \lambda) \cdot (x, y) = (\kappa^2 x, \lambda y)$ and $H = \boldsymbol{\mu}_2$ by $-1 \cdot (x, y) = (-x, -y)$. The box in this example is $\mathcal{B}_{\mathbf{T}} = \{(0, 0), (1, 0)\}$. A rank 1 toric torsion free sheaf $\mathcal{F}$ on $[\mathbb{C}^2/\boldsymbol{\mu}_2]$ corresponds to the following data. Firstly, $\mathcal{F} = {}_{(0,0)}\mathcal{F}$ or $\mathcal{F} = {}_{(1,0)}\mathcal{F}$, because the rank is 1. In either case, ${}_b\mathcal{F}$ is described by specifying a double filtration of $\mathbb{C}$ on $\mathbb{Z}^2$ corresponding to the S-family ${}_bF(\ell_1, \ell_2)$. The "stackiness" comes in by choosing $(L_1, L_2) \in \mathbb{Z}^2$ sufficiently large such that ${}_bF(L_1, L_2) = {}_bF(\infty, \infty) = \mathbb{C}$ and giving this vector space weight 0 or 1 with respect to $X(H) = \mathbb{Z}_2$. Since all maps are the identity (or zero), this fixes all the $X(H)$-weights. A typical ${}_b\hat{F}$ looks like:



where the origin $(\ell_1, \ell_2) = (0, 0)$ can be located anywhere. Here the vector spaces associated to the lattice points on or above the solid line are $\mathbb{C}$ and zero otherwise. Moreover, $0, 1$ refers to the $\boldsymbol{\mu}_2$-weight of the vector space associated to the lattice point in the left-bottom corner of the box containing it.

Example 2. Let $[\mathbb{C}^2/\boldsymbol{\mu}_2]$ be as in the previous example. Let $\mathcal{F}$ be a rank 2 toric torsion free sheaf on $[\mathbb{C}^2/\boldsymbol{\mu}_2]$. Decompose $\mathcal{F} = {}_{(0,0)}\mathcal{F} \oplus {}_{(0,1)}\mathcal{F}$. There are three possibilities:

- (I) The sheaves ${}_{(0,0)}\mathcal{F}$ and ${}_{(0,1)}\mathcal{F}$ are both nonzero: then each is a rank 1 toric torsion free sheaf as in example 1.
- (II) Only one ${}_b\mathcal{F}$ is nonzero, then ${}_bF(L_1, L_2) = {}_bF(\infty, \infty) = \mathbb{C}^2$ for sufficiently large $(L_1, L_2) \in \mathbb{Z}^2$ and this space decomposes as a (nontrivial) sum of a weight 0 and 1 space ${}_bF(L_1, L_2) = {}_bF(L_1, L_2)_0 \oplus {}_bF(L_1, L_2)_1$. The 1-dimensional spaces ${}_bF(L_1, L_2)_n$ both have a double filtration as in example 1 and ${}_b\mathcal{F}$ decomposes as a sum of two rank 1 toric torsion free sheaves.

(III) Only one ${}_b\mathcal{F}$ is nonzero, then ${}_bF(L_1, L_2) = {}_bF(\infty, \infty) = \mathbb{C}^2$ for sufficiently large $(L_1, L_2) \in \mathbb{Z}^2$ and this space has weight 0 or 1. This time ${}_bF(\ell_1, \ell_2)$ is a double filtration of $\mathbb{C}^2$ on $\mathbb{Z}^2$ and the $X(H)$-weight of each ${}_bF(\ell_1, \ell_2)$ is fully determined by the $X(H)$-weight of ${}_bF(L_1, L_2)$.

For a fixed choice of dimensions of the $X(\mathbf{T})$-weight spaces, there are only a finite number of sheaves of type I and II, but the sheaves of type III have continuous moduli corresponding to different choices of inclusions $\mathbb{C} \subset \mathbb{C}^2$. A typical sheaf of type III looks like:

${}_b\hat{F}$



where the vector spaces below the single line are 0-dimensional, on and above but below the double line 1-dimensional and on or above the double line $\mathbb{C}^2$. The 0's and 1's in the diagram denote the $X(H)$-weights. The continuous moduli in this example are $(\mathbb{P}^1)^3$, because there are three connected components between the single and double line.

4. Weighted projective planes

The weighted projective plane $\mathbb{P} := \mathbb{P}(a, b, c)$ is by definition the quotient stack $[\mathbb{C}^3 \backslash \{0\}/\mathbb{C}^*]$, where, denoting by (X, Y, Z) the coordinates on $\mathbb{C}^3$ and $\lambda \in \mathbb{C}^*$ acts by

$$\lambda \cdot (X, Y, Z) = (\lambda^a X, \lambda^b Y, \lambda^c Z).$$

This is a smooth complete toric DM stack in the sense of [BCS, FMN]. Let $d := \gcd(a, b, c)$, $d_{12} := \gcd(a, b)$, $d_{13} := \gcd(a, c)$ and $d_{23} := \gcd(b, c)$. The coarse moduli scheme of $\mathbb{P}$, denoted by $\mathbf{P} = \mathbf{P}(a, b, c)$, is the weighted projective plane in the classical sense [Ful, Sect. 2.2]. The toric variety $\mathbf{P}(a, b, c)$ (in general singular) is isomorphic to $\mathbf{P}(a/d, b/d, c/d)$. If $d_{12} = d_{23} = d_{13} = 1$, then the structure map $\mathbb{P} \to \mathbf{P}$ is an isomorphism away from the points among $(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1) \in \mathbf{P}$ which are singular[6]. In the case $d = 1$, $\mathbb{P}(a, b, c)$ is an orbifold meaning that the structure map $\mathbb{P} \to \mathbf{P}$ is an isomorphism away from the lines $(0 : Y : Z), (X : 0 : Z), (X : Y : 0) \subset \mathbf{P}$. In general, $\mathbb{P}$ is a $B\boldsymbol{\mu}_d$-gerbe over the orbifold $\mathbb{P}(a/d, b/d, c/d)$.

---

[6] In this case $\mathbb{P}(a, b, c)$ is called a canonical DM stack (see [FMN]).

4.1. **Open cover and torus action.** $\mathbb{P}$ can be covered by the standard open substacks $\mathfrak{U}_1, \mathfrak{U}_2, \mathfrak{U}_3$ respectively corresponding to the open sets $\{X \neq 0\}, \{Y \neq 0\}, \{Z \neq 0\} \subset \mathbb{C}^3 \backslash \{0\}$. Define a map of sets

$$\hat{ } : \{1,2,3\} \to \mathbb{Z}_{>0},$$

sending $1 \mapsto a$, $2 \mapsto b$ and $3 \mapsto c$. By [BCS], we have an isomorphism of stacks

$$\mathfrak{U}_i \cong [\mathbb{C}^2 / \boldsymbol{\mu}_{\hat{i}}],$$

for $i = 1,2,3$, where $\mu \in \boldsymbol{\mu}_{\hat{i}}$ acts on $(x,y) \in \mathbb{C}^2$ by

$$\begin{cases} (\mu^b x, \mu^c y) & \text{if } i = 1, \\ (\mu^a x, \mu^c y) & \text{if } i = 2, \\ (\mu^a x, \mu^b y) & \text{if } i = 3. \end{cases}$$

The open immersion $\mathfrak{U}_i \hookrightarrow \mathbb{P}$ is induced by the natural inclusion $\boldsymbol{\mu}_{\hat{i}} \hookrightarrow \mathbb{C}^*$ as the group of $\hat{i}$-th roots of unity and the map $\mathbb{C}^2 \to \mathbb{C}^3 \backslash \{0\}$ sending (x,y) to

$$\begin{cases} (1, x, y) & \text{if } i = 1, \\ (x, 1, y) & \text{if } i = 2, \\ (x, y, 1) & \text{if } i = 3. \end{cases}$$

We compute the double stack theoretic intersections of the $\mathfrak{U}_i$'s by taking the fiber product over $\mathbb{P}$ via the maps defined above. The outcome is

$$\mathfrak{U}_{ij} := \mathfrak{U}_i \times_{\mathbb{P}} \mathfrak{U}_j \cong [\mathbb{C}^* \times \mathbb{C} / \boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}],$$

for $(i,j) = (1,2), (2,3), (1,3)$. The action of $(\mu, \nu) \in \boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}$ on $(\gamma, z) \in \mathbb{C}^* \times \mathbb{C}$ is given by

$$(\nu \mu^{-1} \gamma, \mu^{\hat{k}} z),$$

where $k \in \{1,2,3\} \backslash \{i,j\}$. The open immersions $\mathfrak{U}_{ij} \hookrightarrow \mathfrak{U}_i$ and $\mathfrak{U}_{ij} \hookrightarrow \mathfrak{U}_j$ are induced respectively from the projections

$$\boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}} \to \boldsymbol{\mu}_{\hat{i}} \quad \text{and} \quad \boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}} \to \boldsymbol{\mu}_{\hat{j}},$$

and the maps $\mathbb{C}^* \times \mathbb{C} \to \mathbb{C} \times \mathbb{C}$ that send (γ, z) to respectively

$$\begin{cases} (\gamma^{-b}, z) & \text{if } (i,j) = (1,2), \\ (z, \gamma^{-c}) & \text{if } (i,j) = (2,3), \\ (z, \gamma^{-c}) & \text{if } (i,j) = (1,3), \end{cases} \quad \text{and} \quad \begin{cases} (\gamma^a, z\gamma^c) & \text{if } (i,j) = (1,2), \\ (z\gamma^a, \gamma^b) & \text{if } (i,j) = (2,3), \\ (\gamma^a, z\gamma^b) & \text{if } (i,j) = (1,3). \end{cases}$$

Similarly, one can show

$$\mathfrak{U}_{123} := \mathfrak{U}_1 \times_{\mathbb{P}} \mathfrak{U}_2 \times_{\mathbb{P}} \mathfrak{U}_3 \cong [\mathbb{C}^* \times \mathbb{C}^* / \boldsymbol{\mu}_a \times \boldsymbol{\mu}_b \times \boldsymbol{\mu}_c],$$

where the action of $(\mu, \nu, \xi) \in \boldsymbol{\mu}_a \times \boldsymbol{\mu}_b \times \boldsymbol{\mu}_c$ on $(\kappa, \lambda) \in \mathbb{C}^* \times \mathbb{C}^*$ is given by

$$(\nu\mu^{-1}\kappa, \xi\nu^{-1}\lambda).$$

If (i, j) is one of the pairs

$$(1, 2), (2, 3), (1, 3),$$

then the open immersion $\mathfrak{U}_{123} \hookrightarrow \mathfrak{U}_{ij}$ is given by the corresponding projection

$$\boldsymbol{\mu}_a \times \boldsymbol{\mu}_b \times \boldsymbol{\mu}_c \to \boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}},$$

and the map $\mathbb{C}^* \times \mathbb{C}^* \to \mathbb{C}^* \times \mathbb{C}$ taking (κ, λ) to

$$\begin{cases} (\kappa, \kappa^{-c}\lambda^{-c}) & \text{if } (i, j) = (1, 2), \\ (\lambda, \kappa^a) & \text{if } (i, j) = (2, 3), \\ (\kappa\lambda, \kappa^{-b}) & \text{if } (i, j) = (1, 3). \end{cases}$$

The stack $\mathbb{P}$ is equipped with a natural action of a 2-dimensional torus obtained by rigidifying its open dense substack $\mathfrak{U}_{123} \cong [\mathbb{C}^{*3}/\mathbb{C}^*]$ defined above (see [FMN]). Denote this torus[7] by $\mathbf{T} \cong \mathbb{C}^* \times \mathbb{C}^*$. One can easily define an action of $\mathbf{T}$ on the stacks $\mathfrak{U}_i$ and $\mathfrak{U}_{ij}$ such that all the open immersions defined above are $\mathbf{T}$-equivariant. In the table below we summarize the result by showing the corresponding weights of the $\mathbf{T}$-action on $\mathbb{C} \times \mathbb{C}$ for each $\mathfrak{U}_i$ and on $\mathbb{C}^* \times \mathbb{C}$ for each $\mathfrak{U}_{ij}$ (recall Lemma 2.4):

	$\mathbf{T}$-weights on $\mathbb{C} \times \mathbb{C}$
$\mathfrak{U}_1$	$(b, 0), (0, c)$
$\mathfrak{U}_2$	$(-a, 0), (-c, c)$
$\mathfrak{U}_3$	$(0, -a), (b, -b)$

	$\mathbf{T}$-weights on $\mathbb{C}^* \times \mathbb{C}$
$\mathfrak{U}_{12}$	$(-1, 0), (0, c)$
$\mathfrak{U}_{23}$	$(1, -1), (-a, 0)$
$\mathfrak{U}_{13}$	$(0, -1), (b, 0)$

TABLE 1. $\mathbf{T}$-action on each $\mathfrak{U}_i$ and $\mathfrak{U}_{ij}$.

The $\mathbf{T}$-action on $\mathbb{P}$ has three fixed points P_1, P_2, P_3 which correspond to the respective origins of the open substacks $\mathfrak{U}_1, \mathfrak{U}_2, \mathfrak{U}_3$.

4.2. ***K*-groups of weighted projective planes.** Denote by $K(\mathbb{P})$ be the Grothendieck group of coherent sheaves on $\mathbb{P}$ and let $K(\mathbb{P})_{\mathbb{Q}} := K(\mathbb{P}) \otimes_{\mathbb{Z}} \mathbb{Q}$. By [BH]

$$K(\mathbb{P})_{\mathbb{Q}} \cong \mathbb{Q}[g, g^{-1}]/(1 - g^a)(1 - g^b)(1 - g^c).$$

Here $g = [\mathcal{O}_{\mathbb{P}}(-1)]$ is a generator of $\operatorname{Pic}(\mathbb{P}) \cong \mathbb{Z}$ [FMN, Ex. 7.27]. The classes of the structure sheaves of the fixed points of the $\mathbf{T}$-action are

$$[\mathcal{O}_{P_i}] = (1 - g^a)(1 - g^b)(1 - g^c)/(1 - g^{\hat{i}}).$$

---

[7] In fact $\mathfrak{U}_{abc} \cong \mathbf{T} \times B\boldsymbol{\mu}_d$.

We prove this in Proposition 7.1 as an easy corollary of the general description of toric sheaves on $\mathbb{P}$ developed in Section 5. We get the following identities in $K(\mathbb{P})_{\mathbb{Q}}$

$$[\mathcal{O}_{P_i}] = [\mathcal{O}_{P_i}]g^{\hat{i}}, \quad i = 1,2,3.$$

The elements $[\mathcal{O}_{P_i}]g^j$ for $i = 1,2,3$ and $j = 0, \ldots, \hat{i}-1$ generate the subgroup of $K(\mathbb{P})_{\mathbb{Q}}$ of 0-dimensional coherent sheaves. It is not hard to prove the following relations among these elements:

$$\begin{aligned}
\sum_{i=0}^{a/d-1} [\mathcal{O}_{P_1}]g^{id} &= \sum_{j=0}^{b/d-1} [\mathcal{O}_{P_2}]g^{jd} = \sum_{k=0}^{c/d-1} [\mathcal{O}_{P_3}]g^{kd}, \\
\sum_{i=0}^{a/d-1} [\mathcal{O}_{P_1}]g^{id+1} &= \sum_{j=0}^{b/d-1} [\mathcal{O}_{P_2}]g^{jd+1} = \sum_{k=0}^{c/d-1} [\mathcal{O}_{P_3}]g^{kd+1}, \\
&\cdots \\
\sum_{i=0}^{a/d-1} [\mathcal{O}_{P_1}]g^{id+d-1} &= \sum_{j=0}^{b/d-1} [\mathcal{O}_{P_2}]g^{jd+d-1} = \sum_{k=0}^{c/d-1} [\mathcal{O}_{P_3}]g^{kd+d-1}, \\
\sum_{i=0}^{a/d_{12}-1} [\mathcal{O}_{P_1}]g^{id_{12}} &= \sum_{j=0}^{b/d_{12}-1} [\mathcal{O}_{P_2}]g^{jd_{12}}, \ldots \\
\sum_{j=0}^{b/d_{23}-1} [\mathcal{O}_{P_2}]g^{jd_{23}} &= \sum_{k=0}^{c/d_{23}-1} [\mathcal{O}_{P_3}]g^{kd_{23}}, \ldots \\
\sum_{i=0}^{a/d_{13}-1} [\mathcal{O}_{P_1}]g^{id_{13}} &= \sum_{k=0}^{c/d_{13}-1} [\mathcal{O}_{P_3}]g^{kd_{13}}, \ldots
\end{aligned} \tag{4}$$

For example, the first equality in the first row follows from the following identity in $K(\mathbb{P})_{\mathbb{Q}}$

$$(1 + g^d + \cdots + g^{a-d})(1 - g^b)(1 - g^c) = (1 - g^a)(1 + g^d + \cdots + g^{b-d})(1 - g^c).$$

The first equality in the second row is obtained by multiplying both sides of this equation by g etc.

Let $I\mathbb{P}$ be the inertia stack of $\mathbb{P}$ and $\pi : I\mathbb{P} \to \mathbb{P}$ the natural map. We index the connected components of $I\mathbb{P}$ by the set F and its elements by $f \in F$. For any coherent sheaf $\mathcal{F}$ on $\mathbb{P}$, define

$$\widetilde{\mathrm{ch}} : K(\mathbb{P})_{\mathbb{Q}} \to A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}, \ \widetilde{\mathrm{ch}}(\mathcal{F}) := \sum_{f \in F} \sum_i \omega_{f,i} \cdot \mathrm{ch}(\mathcal{F}_{f,i}),$$

where $\mathcal{F}_f$ is the restriction of $\pi^*\mathcal{F}$ to the component corresponding to f, $\mathcal{F}_f = \bigoplus_i \mathcal{F}_{f,i}$ is its decomposition into eigenvectors and $\omega_{f,i} \in \boldsymbol{\mu}_\infty$ are the corresponding eigenvalues[8]. For any $f \in F$, we denote by $\widetilde{\mathrm{ch}}(\mathcal{F})_f$ the restriction of $\widetilde{\mathrm{ch}}(\mathcal{F})$ to the component Z of $I\mathbb{P}$ corresponding to f.

Define $\widetilde{\mathrm{td}} : \mathrm{Pic}(\mathbb{P}) \to A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ as follows. Let $L \in \mathrm{Pic}(\mathbb{P})$ and let $f \in F$. Denoting $c_1(L_f) = x_f$ and denoting the corresponding eigenvalue of L_f by ω_f, the component $\widetilde{\mathrm{td}}(L)_f$ is given by[9]

$$\frac{x_f^{\delta_{1,\omega_f}}}{1 - \omega_f e^{-x_f}}.$$

The whole point of these definitions is that one can compute the holomorphic Euler characteristic of $\mathcal{F}$ by using Toën-Riemann-Roch (TRR) [Toe]

$$\chi(\mathcal{F}) = \int_{I\mathbb{P}} \widetilde{\mathrm{ch}}(\mathcal{F}) \cdot \widetilde{\mathrm{td}}(\mathcal{O}_{\mathbb{P}}(a)) \cdot \widetilde{\mathrm{td}}(\mathcal{O}_{\mathbb{P}}(b)) \cdot \widetilde{\mathrm{td}}(\mathcal{O}_{\mathbb{P}}(c)).$$

Define $m := \mathrm{lcm}(a, b, c)$. For any coherent sheaf $\mathcal{F}$ on $\mathbb{P}$ and $t \in \mathbb{Z}_{>0}$, define $P(\mathcal{F}, t) := \chi(\mathcal{F} \otimes \mathcal{O}_{\mathbb{P}}(mt))$ to be the Hilbert polynomial of F with respect to[10] $\mathcal{O}_{\mathbb{P}}(m)$.

Proposition 4.1. *For any $r \in \mathbb{Z}$:*

i) if $d \nmid r$ then $P(\mathcal{O}_{\mathbb{P}}(r), t) = 0$,

ii) if $d \mid r$ then $P(\mathcal{O}_{\mathbb{P}}(r), t)$ modulo the t-constant term is given by

$$\frac{dm^2}{2abc}t^2 + m\left(\frac{2r+a+b+c}{2abc}d + \sum_{1 \leq i < j \leq 3} \frac{1}{\hat{i}\hat{j}} \sum_{\substack{h \in \{1, \ldots, d_{ij}-1\} \\ \frac{d_{ij}}{d} \nmid h}} \frac{\omega_{ij}^{hr}}{1 - \omega_{ij}^{h\hat{k}}}\right)t$$

where $\omega_{ij} = e^{2\pi\sqrt{-1}/d_{ij}}$ and $k \in \{1,2,3\} \setminus \{i,j\}$.

Proof. We use the TRR formula above to compute the coefficients of t^2 and t in $\chi(\mathcal{O}_{\mathbb{P}}(r + mt))$. Only 2 and 1-dimensional components of $I\mathbb{P}$ contribute to these terms. Let $D := \{l/d\}_{l=0,\ldots,d-1}$. The 2-dimensional components of $I\mathbb{P}$ are isomorphic to $\mathbb{P}$ and they are in bijection with the elements of D. The eigenvalue of $\mathcal{O}_{\mathbb{P}}(1)$ on $\mathbb{P}_{l/d}$ is $e^{2\pi\sqrt{-1}l/d}$. For $(i,j) = (1,2), (1,3)$ or $(2,3)$, let $D_{ij} := \{l/d_{ij}\}_{l=0,\ldots,d_{ij}-1} \setminus D$. The 1-dimensional components of $I\mathbb{P}$ are all isomorphic to one of $\mathbb{P}(a,b)$, $\mathbb{P}(a,c)$ or $\mathbb{P}(b,c)$. The components isomorphic to $\mathbb{P}(\hat{i},\hat{j})$ are in bijection with the elements of D_{ij}. The eigenvalue of $\mathcal{O}_{\mathbb{P}}(1)$ on $\mathbb{P}(\hat{i},\hat{j})_{l/d_{ij}}$ is $e^{2\pi\sqrt{-1}l/d_{ij}}$. We find the contribution of each component separately. *Contribution of* $\mathbb{P}_{l/d}$*:* The integrand in the TRR formula is

---

[8] Here $\boldsymbol{\mu}_\infty$ is the group of roots of unity. We will only be concerned with the case $\mathcal{F} = L$ is a line bundle, in which case each L_f is an eigensheaf.

[9] Here δ_{1,ω_f} is the Kronecker delta. This expression reduces to the usual Todd class of a line bundle when $\omega_f = 1$.

[10] $\mathcal{O}_{\mathbb{P}}(m)$ is the pull back of $\mathcal{O}_{\mathbf{P}}(1)$ from the coarse moduli space $\mathbf{P}$.

$$e^{2\pi\sqrt{-1}l(r+mt)/d}\cdot e^{(r+mt)x}\cdot\frac{ax}{1-e^{-ax}}\cdot\frac{bx}{1-e^{-bx}}\cdot\frac{cx}{1-e^{-cx}},$$

where x is the first chern class of the pull back of $\mathcal{O}_{\mathbb{P}}(1)$ to this component. The integral over this component modulo the t-constant term is

$$\frac{e^{2\pi\sqrt{-1}lr/d}}{abc}\left((m^2/2)t^2+(r+a/2+b/2+c/2)mt\right).$$

Contribution of $\mathbb{P}(\hat{i},\hat{j})_{l/d_{ij}}$*:* The integrand in the TRR formula is

$$e^{2\pi\sqrt{-1}l(r+mt)/d_{ij}}\cdot e^{(r+mt)x}\cdot\frac{\hat{i}x}{1-e^{-\hat{i}x}}\cdot\frac{\hat{j}x}{1-e^{-\hat{j}x}}\cdot\frac{1}{1-e^{2\pi\sqrt{-1}l\hat{k}/d_{ij}}\cdot e^{-\hat{k}x}},$$

where x is the first Chern class of the pull back of $\mathcal{O}_{\mathbb{P}}(1)$ to this component and $k\in\{1,2,3\}\setminus\{i,j\}$. The integral over this component modulo the t-constant term is

$$\frac{me^{2\pi\sqrt{-1}lr/d_{ij}}}{\hat{i}\hat{j}(1-e^{2\pi\sqrt{-1}l\hat{k}/d_{ij}})}t.$$

The proposition follows by adding the contributions of all the components above. □

Remark 4.2. *Since* $\mathcal{O}_{\mathbb{P}}(m)$ *is the pull back of* $\mathcal{O}_{\mathbf{P}}(1)$ *from the coarse moduli space,* $P(\mathcal{F},t)$ *is equal to the Hilbert polynomial of the push-forward of* $\mathcal{F}$ *to* $\mathbf{P}$ *with respect to* $\mathcal{O}_{\mathbf{P}}(1)$*. If* $d\nmid r$ *then the push forward of* $\mathcal{O}_{\mathbb{P}}(r)$ *to* $\mathbf{P}$ *is zero, which gives another proof of the first part of Proposition 4.1.*

Definition 4.3. In the proof of Proposition 4.1, we defined the sets $D:=\{l/d\}_{l=0,\ldots,d-1}$ and $D_{ij}:=\{l/d_{ij}\}_{l=0,\ldots,d_{ij}-1}\setminus D$ for all $i,j=1,2,3$. In addition, we define for all $i=1,2,3$

$$D_i:=\{l/\hat{i}\}_{l=0,\ldots,\hat{i}-1}\setminus(D\cup D_{ij}\cup D_{ik}),$$

where $j,k\in\{1,2,3\}\setminus\{i\}$ and $j\neq k$. The 2-dimensional components of $I\mathbb{P}$ are all isomorphic to $\mathbb{P}$ and they are indexed by D. The 1-dimensional components of $I\mathbb{P}$ are isomorphic to $\mathbb{P}(\hat{i},\hat{j})$ and are index by D_{ij} for any $i,j=1,2,3$. Finally, the 0-dimensional components of $I\mathbb{P}$ are isomorphic to $\mathbb{P}(\hat{i})$ and are indexed by D_i, where $i=1,2,3$. The sets D, D_i's and D_{ij}'s give a partition of the set of connected components F of $I\mathbb{P}$. The eigenvalue of $\mathcal{O}_{\mathbb{P}}(1)$ pulled back to $I\mathbb{P}$ and restricted to the component corresponding to $f\in F$ is $e^{2\pi\sqrt{-1}f}$. ⊘

Corollary 4.4. *Let* $\mathcal{E} = \oplus_{u=0}^{E-1} \mathcal{O}_{\mathbb{P}}(u)$ *and assume* $d_{ij} \mid E$ *for all* $1 \leq i < j \leq 3$. *Then for any* $r \in \mathbb{Z}$, *the Hilbert polynomial* $P(\mathcal{O}_{\mathbb{P}}(r) \otimes \mathcal{E}, t)$ *modulo* t*-constant term is given by*

$$P(\mathcal{O}_{\mathbb{P}}(r) \otimes \mathcal{E}, t) = \frac{Em^2}{2abc} t^2 + \sum_{i=1}^{e} \frac{2r + 2u_i + a + b + c}{2abc} mdt,$$

where[11] $u_1, \ldots, u_e$ *are those* $u \in \{0, \ldots, E-1\}$ *for which* $d \mid r + u$.

5. Toric sheaves on weighted projective planes

The toric DM stack $\mathbb{P} := \mathbb{P}(a,b,c)$ has an action of an algebraic torus $\mathbf{T} = \mathbb{C}^{*2}$ as described in Section 4.1. In this section, we describe toric sheaves on $\mathbb{P}$. The approach is as follows. We cover $\mathbb{P}$ by the $\mathbf{T}$-invariant open substacks $\mathfrak{U}_i \cong [\mathbb{C}^2 / \boldsymbol{\mu}_{\hat{i}}]$ for $i = 1, 2, 3$ (see Section 4.1 for the definition of the $\mathfrak{U}_i$'s and $\hat{i}$). The action of $\mathbf{T}$ and $\boldsymbol{\mu}_{\hat{i}}$ on $\mathbb{C}^2$ are described explicitly in Section 4.1. A toric sheaf $\mathcal{F}$ on $\mathbb{P}$ restricts to a toric sheaf $\mathcal{F}_i$ on $\mathfrak{U}_i$ which is (up to equivariant isomorphism) determined by its stacky S-family $\hat{F}_i$ (Proposition 3.5). Conversely, given toric sheaves $\mathcal{F}_i$ on $\mathfrak{U}_i$, we want to find gluing conditions on the stacky S-families $\hat{F}_i$ allowing us to glue the $\mathcal{F}_i$ to a toric sheaf $\mathcal{F}$ on $\mathbb{P}$.

Let $\mathcal{F}_i|_{ij}$ be the pull-back of $\mathcal{F}_i$ along the inclusion $\mathfrak{U}_{ij} = \mathfrak{U}_i \times_{\mathbb{P}} \mathfrak{U}_j \hookrightarrow \mathfrak{U}_i$. In section 4.1, we described these maps explicitly at the groupoid level and we noted $\mathfrak{U}_{ij} \cong [\mathbb{C}^* \times \mathbb{C} / \boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}]$. This will allows us to compute the stacky S-family $\hat{F}_{i,ij}$ of $\mathcal{F}_i|_{ij}$. Matching $\hat{F}_{i,ij}$ and $\hat{F}_{j,ij}$ for all i, j allows us to glue sheaves (Proposition 2.2).

5.1. **Graded tensor products.** Given a morphism of affine schemes $f :$ Spec $S \to$ Spec R and a quasi-coherent sheaf $\mathcal{F}$ corresponding to an R-module $H^0(\mathcal{F})$ on Spec R, the pull-back $f^*\mathcal{F}$ corresponds to the S-module $H^0(f^*\mathcal{F}) \cong H^0(\mathcal{F}) \otimes_R S$ [Har, Prop. 5.2]. Suppose the affine schemes have an action of an algebraic torus $\mathbf{T}$, f is $\mathbf{T}$-equivariant and $\mathcal{F}$ is a toric sheaf. Then $f^*\mathcal{F}$ is a toric sheaf. The $\mathbf{T}$-equivariant structure of $\mathcal{F}$ corresponds to an $X(\mathbf{T})$-grading on $H^0(\mathcal{F})$ and the $\mathbf{T}$-equivariant structure of $f^*\mathcal{F}$ corresponds to an $X(\mathbf{T})$-grading on $H^0(f^*\mathcal{F})$. How can this latter grading be described in terms of the gradings of $H^0(\mathcal{F})$, R and S? We follow Perling's discussion [Per1, Sect. 3].

Suppose we are given, in general, an abelian group A, an A-graded ring R and A-graded modules M and N. Then $M \otimes_R N$ has a natural A-grading as follows [Per1, Sect. 3]. The degree a part $(M \otimes_R N)_a$ is the abelian subgroup generated by all elements

---

[11] In particular, if $d = 1$ then $e = E = \operatorname{rank} \mathcal{E}$.

$$m \otimes n, \text{ with } m \in M_{a'} \text{ and } n \in N_{a''} \text{ such that } a' + a'' = a.$$

Next, suppose we are given an A-graded ring R and a B-graded ring S. A morphism of graded rings is a pair (ϕ, ψ) with $\phi : R \to S$ a ring homomorphism and $\psi : A \to B$ a group homomorphism satisfying $\phi(R_a) \subset S_{\psi(a)}$. Given such a morphism of graded rings, R can be given a B-grading by defining

$$R_b := \bigoplus_{a \in \psi^{-1}(b)} R_a.$$

Likewise, any A-graded R-module M can be a seen as a B-graded R-module.

Back to the geometric setting, the $\mathbf{T}$-equivariant morphism $f : \operatorname{Spec} S \to \operatorname{Spec} R$ gives a morphism of graded rings $(\phi, \chi) : (R, X(\mathbf{T})) \to (S, X(\mathbf{T}))$. We can see $H^0(\mathcal{F})$ and S as R-modules with an $X(\mathbf{T})$-grading. This makes $H^0(f^*\mathcal{F})$ into an $X(\mathbf{T})$-graded S-module. *Note*: in general the $X(\mathbf{T})$-grading on $H^0(\mathcal{F})$ and R is induced by χ, which need not be the identity map. In our applications χ is always the identity map.

5.2. **Gluing conditions.** We are now ready to describe toric sheaves on $\mathbb{P}$. A toric sheaf $\mathcal{F}_i$ on $\mathfrak{U}_i \cong [\mathbb{C}^2/\boldsymbol{\mu}_{\hat{i}}]$ is determined (up to equivariant isomorphism) by its stacky S-family $\hat{F}_i$ (Proposition 3.5). We want to compute the stacky S-family $\hat{F}_{i,ij}$ of the toric sheaf $\mathcal{F}_i|_{ij}$ on $\mathfrak{U}_{ij} \cong [\mathbb{C}^* \times \mathbb{C}/\boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}]$ for all i, j. Matching $\hat{F}_{i,ij}$ and $\hat{F}_{j,ij}$ for all i, j allows us to glue the $\mathcal{F}_i$ to a global toric sheaf $\mathcal{F}$ by Proposition 2.2.

The torus $\mathbf{T}$ has character group $X(\mathbf{T}) = \mathbb{Z}^2$. This torus is acting on each $\mathfrak{U}_i$ and $\mathfrak{U}_{ij}$. We denote the elements of $\mathbf{T} = \mathbb{C}^{*2}$ by $(\kappa, \lambda) \in \mathbf{T}$. For each $\mathfrak{U}_i \cong [\mathbb{C}^2/\boldsymbol{\mu}_{\hat{i}}]$, there are coordinates (x, y) on $\mathbb{C}^2$ such that $(\kappa, \lambda) \cdot (x, y) = (\chi(m_1)(\kappa, \lambda)x, \chi(m_2)(\kappa, \lambda)y)$ for independent $m_1, m_2 \in X(\mathbf{T})$. The characters m_1, m_2 are unique up to *order* and we choose the following orders[12] (see Table 4.1 in Section 4.1)

$$\begin{aligned} i = 1 &: \ (m_1, m_2) = ((b, 0), (0, c)), \\ i = 2 &: \ (m_1, m_2) = ((-c, c), (-a, 0)), \\ i = 3 &: \ (m_1, m_2) = ((0, -a), (b, -b)). \end{aligned}$$

These vectors span the box $\mathcal{B}_{\mathbf{T}}$ of the action of $\mathbf{T}$ on $\mathfrak{U}_i$. Likewise, for each $\mathfrak{U}_{ij} \cong [\mathbb{C}^* \times \mathbb{C}/\boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}]$ there are coordinates (γ, z) on $\mathbb{C}^* \times \mathbb{C}$ such that $(\kappa, \lambda) \cdot (\gamma, z) = (\chi(m_1)(\kappa, \lambda)\gamma, \chi(m_2)(\kappa, \lambda)z)$ for independent $m_1, m_2 \in X(\mathbf{T})$.

---

[12] Compared to Table 4.1 in Section 4.1, we have changed the order for $i = 2$ because of symmetry reasons.

We compute $\hat{F}_{1,12}$ in terms of $\hat{F}_1$. Recall that we denote the various vector spaces corresponding to $\hat{F}_1$ by ${}_{(i/b,j/c)}F_1(\ell_1,\ell_2)_l$ (see Section 3). Here the box $\mathcal{B}_T = [0,b-1]\times[0,c-1] \subset X(\mathbf{T})$ and its elements are written by $(i/b,j/c) \in \mathcal{B}_T$. The inclusion $\mathfrak{U}_{12} \cong [\mathbb{C}^* \times \mathbb{C}/\boldsymbol{\mu}_a \times \boldsymbol{\mu}_b] \hookrightarrow \mathfrak{U}_1 \cong [\mathbb{C}^2/\boldsymbol{\mu}_a]$ corresponds to the étale morphism (Section 4.1)

$$\mathbb{C}^* \times \mathbb{C} \longrightarrow \mathbb{C}^2,\ (\gamma,z) \mapsto (\gamma^{-b},z) = (x,y), \tag{5}$$

which intertwines the $\boldsymbol{\mu}_a \times \boldsymbol{\mu}_b$- and $\boldsymbol{\mu}_a$-actions and the $\mathbf{T}$-actions. These actions are given by

$$\begin{aligned}(\kappa,\lambda)\cdot(\gamma,z) &= (\kappa^{-1}\gamma,\lambda^c z),\\ (\mu,\nu)\cdot(\gamma,z) &= (\mu^{-1}\nu\gamma,\mu^c z),\\ (\kappa,\lambda)\cdot(x,y) &= (\kappa^b x,\lambda^c y),\\ \mu\cdot(x,y) &= (\mu^b x,\mu^c y).\end{aligned}$$

We denote a-th roots of unity by $\mu \in \boldsymbol{\mu}_a$, b-th roots of unite by $\nu \in \boldsymbol{\mu}_b$ and c-th roots of unity by $\xi \in \boldsymbol{\mu}_c$. We first restrict $\mathcal{F}_1$ to $\mathbb{C}^* \times \mathbb{C} \subset \mathbb{C}^2$. For a fixed ℓ_2, the vector spaces ${}_{(i/b,j/c)}F_1(\ell_1,\ell_2)$ stabilize for $\ell_1 \gg 0$. This means they are all isomorphic for $\ell_1 \gg 0$. Up to isomorphism of $\hat{F}_1$, we can take all these isomorphisms between these limiting spaces to be the identity map (though these maps in general do swap the fine-grading!). We denote the limit by ${}_{(i/b,j/c)}F_1(\infty,\ell_2)$. The sheaf $\mathcal{F}_1|_{\mathbb{C}^*\times\mathbb{C}}$ has an S-family $\hat{G}_1$ which is easily computed using the description of grading on a tensor product of Section 5.1

$${}_{(i/b,j/c)}G_1(\ell_1,\ell_2) = {}_{(i/b,j/c)}F_1(\infty,\ell_2).$$

The maps between these vector spaces are the identity in the ℓ_1-direction and the obvious induced maps in the ℓ_2-direction. This is because multiplication by x is an isomorphism. The fine-grading does depend on ℓ_1 albeit in a trivial way. This means that for $\ell_1 \gg 0$ such that ${}_{(i/b,j/c)}F_1(\ell_1,\ell_2) = {}_{(i/b,j/c)}F_1(\infty,\ell_2)$, we have ${}_{(i/b,j/c)}G_1(\ell_1,\ell_2)_l = {}_{(i/b,j/c)}F_1(\ell_1,\ell_2)_l$. The fine-grading on ${}_{(i/b,j/c)}G_1(\ell_1,\ell_2)$ for one fixed ℓ_1 determines the the fine-grading on ${}_{(i/b,j/c)}G_1(\ell_1,\ell_2)$ for all ℓ_1

$${}_{(i/b,j/c)}G_1(\ell_1,\ell_2)_l = {}_{(i/b,j/c)}G_1(0,\ell_2)_{l-\ell_1 b} \otimes \mu^{\ell_1 b}.$$

Here $-\otimes\mu^l$ stands for tensoring with the one-dimensional representation of $\boldsymbol{\mu}_a$ of weight $l \in \mathbb{Z}_a$. We *define* the fine-grading on the limit vector space ${}_{(i/b,j/c)}F_1(\infty,\ell_2)$ by

$${}_{(i/b,j/c)}F_1(\infty,\ell_2)_l := {}_{(i/b,j/c)}G_1(0,\ell_2)_l.$$

The S-family $\hat{F}_{1,12}$ at the point $(i_0/b, j/c) + \ell_1 \cdot (b,0) + \ell_2 \cdot (0,c) \in X(\mathbf{T}) = \mathbb{Z}^2$ is

$$\bigoplus_{0 \leq i \leq b-1} {}_{(i/b,j/c)}F_1(\infty, \ell_2).$$

The maps between these vector spaces are the identity in the ℓ_1-direction and the obvious induced maps in the ℓ_2-direction. This expression can be derived as follows. Using the description of grading on a tensor product of Section 5.1 and the explicit formula for the map (5), we see that an element of $\hat{F}_{1,12}$ at $(i_0/b, j/c) + \ell_1 \cdot (b,0) + \ell_2 \cdot (0,c) \in X(\mathbf{T}) = \mathbb{Z}^2$ can be uniquely written in the form

$$\bigoplus_{0 \leq i \leq b-1} v_i \otimes \gamma^{i-i_0},$$

where

$$v_i \in {}_{(i/b,j/c)}G_1(\ell_1, \ell_2).$$

Here the notation $- \otimes \gamma$ comes from the following. The module $F_1 = H^0(\mathcal{F}_1)$ is a finitely generated $\mathbb{C}[x,y]$-module. First, it gets restricted to $\mathbb{C}^* \times \mathbb{C}$, which makes it into a $\mathbb{C}[x^{\pm}, y]$-module G_1. Then it gets pulled-back along the étale morphism (5), which gives us a $\mathbb{C}[\gamma^{\pm}, z]$-module $G_1 \otimes \mathbb{C}[\gamma^{\pm}, z]$. The various homogeneous elements $v_i \otimes \gamma^{i-i_0}$ are elements of this module.

Next we determine the fine-grading. Assuming v_i to be homogenous of weight $l + i - i_0 \in \mathbb{Z}_a$ with respect to the action of μ_a, the weight of $v_i \otimes \gamma^{i-i_0}$ with respect to the action of $\boldsymbol{\mu}_a \times \boldsymbol{\mu}_b$ is

$$(l, i - i_0) \in \mathbb{Z}_a \oplus \mathbb{Z}_b.$$

Therefore, the fine-grading on the weight space of $\hat{F}_{1,12}$ at the point $(i_0/b, j/c) + \ell_1 \cdot (b,0) + \ell_2 \cdot (0,c) \in X(\mathbf{T}) = \mathbb{Z}^2$ is

$$\bigoplus_{\substack{0 \leq i \leq b-1 \\ l \in \mathbb{Z}_a}} {}_{(i/b,j/c)}G_1(\ell_1, \ell_2)_{l+i-i_0} \otimes \mu^{i_0 - i} \otimes \nu^{i-i_0},$$

or in other words

$$\bigoplus_{\substack{0 \leq i \leq b-1 \\ l \in \mathbb{Z}_a}} {}_{(i/b,j/c)}F_1(\infty, \ell_2)_{l+i-i_0} \otimes \mu^{i_0 - i + \ell_1 b} \otimes \nu^{i-i_0}.$$

Here $- \otimes \mu^l$ stands for tensoring with the one-dimensional representation of $\boldsymbol{\mu}_a$ of weight $l \in \mathbb{Z}_a$ and $- \otimes \nu^l$ stands for tensoring with the one-dimensional representation of $\boldsymbol{\mu}_b$ of weight $l \in \mathbb{Z}_b$.

Theorem 5.1 (Gluing conditions)**.** *The category of toric sheaves on* $\mathbb{P}$ *is equivalent to the category of triples* $\{\hat{F}_i\}_{i=1,2,3}$ *of stacky finite* S*-families on* $\mathfrak{U}_i \cong [\mathbb{C}^2/\boldsymbol{\mu}_{\hat{i}}]$ *satisfying the following gluing conditions:*

$$\bigoplus_{\substack{0 \leq i \leq b-1 \\ l \in \mathbb{Z}_a}} {}_{(i/b,j/c)}F_1(\infty,\ell)_l \otimes \mu^{-i} \otimes \nu^i = \bigoplus_{\substack{0 \leq i \leq a-1 \\ l \in \mathbb{Z}_b}} {}_{(j/c,i/a)}F_2(\ell,\infty)_l \otimes \nu^{-i-j-\ell c} \otimes \mu^{i+j+\ell c},$$

$$\bigoplus_{\substack{0 \leq i \leq c-1 \\ l \in \mathbb{Z}_b}} {}_{(i/c,j/a)}F_2(\infty,\ell)_l \otimes \nu^{-i} \otimes \xi^i = \bigoplus_{\substack{0 \leq i \leq b-1 \\ l \in \mathbb{Z}_c}} {}_{(j/a,i/b)}F_3(\ell,\infty)_l \otimes \xi^{-i-j-\ell a} \otimes \nu^{i+j+\ell a},$$

$$\bigoplus_{\substack{0 \leq i \leq a-1 \\ l \in \mathbb{Z}_c}} {}_{(i/a,j/b)}F_3(\infty,\ell)_l \otimes \xi^{-i} \otimes \mu^i = \bigoplus_{\substack{0 \leq i \leq c-1 \\ l \in \mathbb{Z}_a}} {}_{(j/b,i/c)}F_1(\ell,\infty)_l \otimes \mu^{-i-j-\ell b} \otimes \xi^{i+j+\ell b},$$

for all $\ell \in \mathbb{Z}$ *and similar gluing conditions between the linear maps.*

Proof. The category of toric sheaves on $\mathbb{P}$ is equivalent to the category of triples $\{\mathcal{F}_i\}_{i=1,2,3}$ of toric sheaves on $\mathfrak{U}_i \cong [\mathbb{C}^2/\boldsymbol{\mu}_{\hat{i}}]$ satisfying the gluing conditions of Proposition 2.2. In turn, a toric sheaf $\mathcal{F}_i$ is described (up to equivariant isomorphism) by its stacky S-family $\hat{F}_i$ (Proposition 3.5). Given toric sheaves $\{\mathcal{F}_i\}_{i=1,2,3}$ on $\mathfrak{U}_i \cong [\mathbb{C}^2/\boldsymbol{\mu}_{\hat{i}}]$, we consider the toric sheaves $\mathcal{F}_i|_{ij}$ on $\mathfrak{U}_{ij} \cong [\mathbb{C}^* \times \mathbb{C}/\boldsymbol{\mu}_{\hat{i}} \times \boldsymbol{\mu}_{\hat{j}}]$ for all i,j. We compute the stacky S-family $\hat{F}_{i,ij}$ corresponding to $\mathcal{F}_i|_{ij}$.

The case $\hat{F}_{1,12}$ was computed. At a point $(0, j + c\ell) \in X(\mathbf{T}) = \mathbb{Z}^2$ its stacky S-family is given by

$$\bigoplus_{\substack{0 \leq i \leq b-1 \\ l \in \mathbb{Z}_a}} {}_{(i/b,j/c)}F_1(\infty,\ell)_l \otimes \mu^{-i} \otimes \nu^i.$$

Completely similarly, the stacky S-family of $\hat{F}_{2,12}$ at $(-j-\ell c, j+\ell c) \in X(\mathbf{T}) = \mathbb{Z}^2$ is given by

$$\bigoplus_{\substack{0 \leq i \leq a-1 \\ l \in \mathbb{Z}_b}} {}_{(j/c,i/a)}F_2(\ell,\infty)_l \otimes \nu^{-i} \otimes \mu^i.$$

Note however, that we cannot "equate" these two answers. This is because the first vector spaces live on the line $(0, j + c\ell)$ and the second vector spaces live on the line $(-j - \ell c, j + \ell c) \in X(\mathbf{T}) = \mathbb{Z}^2$. Translating these last vector spaces to the line $(0, j + \ell c) \in X(\mathbf{T}) = \mathbb{Z}^2$ gives the first equation. The other equations go similar.

Note that the equalities in the theorem should be isomorphisms of graded vector spaces satisfying the cocycle identity. However, up to isomorphism (and hence up to equivalence of categories), we can choose them to be identities. □

6. Examples of toric sheaves on weighted projective planes

In this section, we give some examples of toric sheaves on $\mathbb{P} := \mathbb{P}(a,b,c)$ as described by Theorem 5.1. The focus will be on rank 1 and 2 toric torsion free sheaves and vector bundles.

Example 1. Let $\mathcal{F}$ be a rank 1 toric torsion free sheaf on $\mathbb{P}^2$. In each chart $\mathfrak{U}_i \cong \mathbb{C}^2$, $\mathcal{F}$ is described by its S-family $\hat{F}_i$ (Section 3)



where on and above the solid line all weight spaces are $\mathbb{C}$ and all maps are the identity and below the solid line all weight spaces and maps are 0. The solid dot indicates an element $(A_j, A_k) \in \mathbb{Z}^2$ which determines the position of the partition. According to Theorem 5.1, this glues to a toric sheaf if and only if

$$A_2 = A_3,\ A_4 = A_5,\ A_6 = A_1.$$

Example 2. Let $\mathcal{F}$ be a rank 1 toric torsion free sheaf on $\mathbb{P}(1,1,2)$. The charts $\mathfrak{U}_1$ and $\mathfrak{U}_2$ have a box of size 2, whereas $\mathfrak{U}_3$ has a trivial box. Since we are in the rank 1 case, for each $i = 1,2,3$ there is only one $b_i \in \mathcal{B}_{\mathbf{T}}$ for which the stacky S-family ${}_{b_i}\hat{F}_i$ is nonzero (Section 3). A priori this gives $2 \cdot 2 = 4$ choices of b_i's. The gluing conditions reduce this to 2 choices

$$b_1 = (0, i/2),\ b_2 = (i/2, 0),\ b_3 = (0,0),\ i \in \{0,1\}.$$

Fix one of these 2 choices. On each chart $\mathfrak{U}_i$, the S-family ${}_{b_i}\hat{F}_i$ is given by the same data as in the previous example. Here we temporarily ignore the fact that the weight spaces in $\mathfrak{U}_3$ carry a $\boldsymbol{\mu}_2$-action. This local data is described by 3 partitions and 6 integers $A_1, \ldots, A_6 \in \mathbb{Z}$. The gluing conditions impose $A_2 = A_3,\ A_4 = A_5,\ A_6 = A_1$ as in the previous example. On the chart $\mathfrak{U}_3$, each weight space carries a $\boldsymbol{\mu}_2$-action. Fixing the weight of one nonzero weight space determines the weights of all other weight spaces. Since $\boldsymbol{\mu}_2$ acts by $(-1,-1)$, the weights alternate as in example 1 of Section 3. For fixed partitions and integers A_i, this a priori gives 2 choices of $\boldsymbol{\mu}_2$-actions. However, the gluing condition allow precisely one choice! (It is immediate there can be at most one choice. Looking closer one finds that there is a choice.)

Example 3. Let $\mathcal{F}$ be a rank 1 toric torsion free sheaf on $\mathbb{P}(2,2,2)$. This time, each chart $\mathfrak{U}_i$ has a box of size $2 \cdot 2 = 4$. As before, for each $i = 1,2,3$ there is only one $b_i \in \mathcal{B}_{\mathbf{T}}$ for which ${}_{b_i}\hat{F}_i$ is nonzero (Section 3). A priori this gives $4 \cdot 4 \cdot 4 = 64$ choices of b_i's. The gluing conditions reduce this to 8 choices

$$b_1 = (i/2, j/2),\ b_2 = (j/2, k/2),\ b_3 = (k/2, i/2),\ i,j,k \in \{0,1\}.$$

Fix any of these 8 choices. On each chart $\mathfrak{U}_i$, the S-family ${}_{b_i}\hat{F}_i$ is given by the same data as in the previous two examples (again, first ignoring the $\boldsymbol{\mu}_2$-actions). This local data is described by 3 partitions and 6 integers $A_1, \ldots, A_6 \in \mathbb{Z}$. The gluing conditions impose $A_2 = A_3,\ A_4 = A_5,\ A_6 = A_1$ as before. On each chart $\mathfrak{U}_i$, the weight spaces carry a $\boldsymbol{\mu}_2$-action. Since the finite groups $\boldsymbol{\mu}_2$ act trivially, all weight spaces on a fixed chart $\mathfrak{U}_i$ have the same character. For fixed partitions and integers A_i, this a priori gives $2 \cdot 2 \cdot 2 = 8$ choices of $\boldsymbol{\mu}_2$-actions. However, the gluing condition again allow precisely one choice!

From examples 1,2 and 3 the general picture of rank 1 toric torsion free sheaves on any $\mathbb{P}$ is easily obtained. We start by describing toric line bundles. A toric line bundle L is locally described by three stacky S-families $\hat{L}_i$ and for each $i = 1,2,3$ there is only one $b_i \in \mathcal{B}_{\mathbf{T}}$ for which ${}_{b_i}\hat{L}_i$ is nonzero. A priori this gives $bc + ac + ba$ choices, but the gluing conditions reduce this to one of the following abc choices

$$\begin{aligned} &b_1 = (i/b, j/c),\ b_2 = (j/c, k/a),\ b_3 = (k/a, i/b), \\ &0 \leq i \leq b-1,\ 0 \leq j \leq c-1,\ 0 \leq k \leq a-1. \end{aligned}$$

Fix any such choice i,j,k. On each chart $\mathfrak{U}_i$, the S-family ${}_{b_i}\hat{L}_i$ is fully determined by the weight of its unique homogeneous generator, i.e. by two integers. Here we ignore the $\boldsymbol{\mu}_{\hat{i}}$-actions for the moment. A priori this gives 6 integers to choose, but the gluing conditions reduce this to 3. In particular, there are unique integers[13] I, J, K such that the homogeneous generators have torus weight

$$\begin{aligned} i/b(b,0) + j/c(0,c) + I(b,0) + J(0,c) &\in X(\mathbf{T}) = \mathbb{Z}^2, \\ j/c(-c,c) + k/a(-a,0) + J(-c,c) + K(-a,0) &\in X(\mathbf{T}) = \mathbb{Z}^2, \\ k/a(0,-a) + i/b(b,-b) + K(0,-a) + I(b,-b) &\in X(\mathbf{T}) = \mathbb{Z}^2. \end{aligned}$$

---

[13] In Examples 1–3 of this section I, J, K are respectively denoted by A_1, A_3, A_5.

Defining $A = i + Ib$, $B = j + Jc$ and $C = k + Ka$, we see that the data so far is uniquely determined by the specification of three integers A, B, C. In other words, specifying $A, B, C \in \mathbb{Z}$ determines the positions of the homogeneous generators in the charts $\mathfrak{U}_1, \mathfrak{U}_2, \mathfrak{U}_3$ according to the above recipe. So far we have ignored the $\boldsymbol{\mu}_{\hat{i}}$-actions. For fixed $A, B, C \in \mathbb{Z}$, there are unique finite group weights of the homogeneous generators for which the data glues to a toric line bundle, i.e. (Theorem 5.1)

$$\begin{aligned} A + B + C &\in \mathbb{Z}_a \text{ on } \mathfrak{U}_1, \\ A + B + C &\in \mathbb{Z}_b \text{ on } \mathfrak{U}_2, \\ A + B + C &\in \mathbb{Z}_c \text{ on } \mathfrak{U}_3. \end{aligned}$$

We denote the toric line bundle corresponding to $(A, B, C) \in \mathbb{Z}^3$ by $L_{(A,B,C)}$ and its stacky S-families by $\hat{L}_{(A,B,C),i}$. From the description of grading on a tensor product (Section 5.1), it is easy to see that

$$L_{(A,B,C)} \otimes L_{(A',B',C')} \cong L_{(A+A',B+B',C+C')}.$$

Proposition 6.1. *Let* $\mathrm{Pic}^{\mathbf{T}}(\mathbb{P})$ *be the equivariant Picard group of* $\mathbb{P}$. *Then*

$$(A, B, C) \in \mathbb{Z}^3 \mapsto L_{(A,B,C)} \in \mathrm{Pic}^{\mathbf{T}}(\mathbb{P})$$

is a group isomorphism.

Recall that the ordinary Picard group of $\mathbb{P}$ is $\mathbb{Z}$ [FMN, Ex. 7.27]. A rank 1 toric torsion free sheaf on $\mathbb{P}$ is described by the same data as a toric line bundle on $\mathbb{P}$ except that on each chart $\mathfrak{U}_i$ a partition is cut out in the corner as in examples 1,2 and 3. We now turn to rank 2.

Example 4. Let $\mathcal{F}$ be a rank 2 toric torsion free sheaf on $\mathbb{P}^2$ (see also [Kly2] or [Koo2]). In each chart $\mathfrak{U}_i \cong \mathbb{C}^2$, $\mathcal{F}$ is described by a double filtration of $\mathbb{C}^2$ (Section 3):



Here $A_1, \ldots, A_6 \in \mathbb{Z}$ fix the position of the filtrations and $\Delta_1, \ldots, \Delta_6 \in \mathbb{Z}_{\geq 0}$ are the widths of the indicated regions. Furthermore, $p_1, \ldots, p_6, q_1, q_2 \subset \mathbb{C}^2$

are 1-dimensional linear subspaces. In each region, the vector spaces on the associated lattice points are as indicated and all maps are inclusions. Also, the lattice points below the single solid line have 0 as their associated vector space. According to Theorem 5.1, this glues to a toric sheaf if and only if

$$\begin{aligned}&A_2 = A_3,\ A_4 = A_5,\ A_6 = A_1,\\ &\Delta_2 = \Delta_3,\ \Delta_4 = \Delta_5,\ \Delta_6 = \Delta_1,\\ &p_2 = p_3 \subset \mathbb{C}^2,\ p_4 = p_5 \subset \mathbb{C}^2,\ p_6 = p_1 \subset \mathbb{C}^2.\end{aligned}$$

Note that the vector spaces q_1, q_2 are not involved in the gluing conditions. A general rank 2 toric torsion free sheaf can have many more (or none) q_i's sitting in the corners. Also, the widths Δ_i are allowed to be zero in which case the corresponding p_i's do not appear in the gluing conditions. Rank 2 toric vector bundles are fully determined by integers $A_1, A_2, A_3 \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ and a point $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$ (Section 3, Equation (3)). Abbreviating $p_{ij} := p_i \cap p_j \subset \mathbb{C}^2$, a typical rank 2 toric vector bundle looks like:







Here the regions labelled by p_{ij} either have associated vector space 0 (if $p_i \neq p_j$) or $p_i = p_j \subset \mathbb{C}^2$ on the lattice points. (So the wiggly line is either not present or a single solid line.) Moreover, the widths Δ_i are allowed to be zero, in which case the corresponding p_i's do not appear.

Example 5. Let $\mathcal{F}$ be a rank 2 toric torsion free sheaf on $\mathbb{P}(1,1,2)$. We focus on the case $\mathcal{F}$ is a vector bundle. The general case can be deduced by the reader. The stacky S-families decompose according to the box elements $\hat{F}_1 = {}_{(0,0)}\hat{F}_1 \oplus {}_{(0,1/2)}\hat{F}_1$, $\hat{F}_2 = {}_{(0,0)}\hat{F}_2 \oplus {}_{(1/2,0)}\hat{F}_2$ and $\hat{F}_3 = {}_{(0,0)}\hat{F}_3$. Since the rank is 2, either each summand is a rank 1 toric line bundle or one summand is zero and the other is a rank 2 toric vector bundle. We distinguish the following cases:

(I) One box summand of $\hat{F}_i$ is zero for both $i = 1, 2$. Let ${}_{b_i}\hat{F}_i \neq 0$ then

$$b_1 = (0, i/2),\ b_2 = (i/2, 0),\ b_3 = (0,0),\ i \in \{0,1\},$$

so there are 2 choices. Fix the b_i's. Disregarding the $\boldsymbol{\mu}_2$-actions for the moment, the S-families $\hat{F}_i$ are exactly as in the case of toric vector bundles in the previous example, i.e. determined by integers $A_1, A_2, A_3 \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ and a point $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$. Again, the widths Δ_i are allowed to be zero, in which case the corresponding p_i's do not appear. For any such fixed choice, there is a unique choice of $\boldsymbol{\mu}_2$-actions on the weight spaces of $\hat{F}_3$ making this into a stacky S-family. For this choice, the limiting vector spaces $\mathbb{C}^2$ do not decompose as the sum of 1-dimensional spaces with different weights.

(II) None of the ${}_{b_i}\hat{F}_i$ is zero. Start by ignoring the $\boldsymbol{\mu}_2$-actions. The S-families ${}_{(0,0)}\hat{F}_1$ and ${}_{(0,0)}\hat{F}_2$ look like





where p is a 1-dimensional vector space and $A_1, A_2, A_3 \in \mathbb{Z}$. Here we have already glued ${}_{(0,0)}\hat{F}_1$ and ${}_{(0,0)}\hat{F}_2$ along $\mathfrak{U}_{12}$. A similar picture can be drawn for ${}_{(0,1/2)}\hat{F}_1$ and ${}_{(1/2,0)}\hat{F}_2$ replacing p, A_i by q, B_i. It is easy to work out what the rest of the gluing conditions for this vector bundle are. We do one case. Assume $A_1 < B_1$, $A_2 > B_2$ and $A_3 > B_3$. In order to satisfy the gluing conditions (still ignoring the $\boldsymbol{\mu}_2$-actions), the vector spaces p, q must be distinct linear subspaces of $\mathbb{C}^2$ (so $\mathbb{C}^2 = p \oplus q$) and the S-family $\hat{F}_3$ must look as follows:



There is a unique choice of $\boldsymbol{\mu}_2$-actions on the weight spaces which makes this into a rank 2 toric vector bundle. Such toric vector bundles are always equivariantly *decomposable* as a sum of two toric line bundles. Even when ignoring the $\boldsymbol{\mu}_2$-actions, there is no analog to this kind of toric vector bundle on $\mathbb{P}^2$. All other type II cases go similar and are also equivariantly decomposable as a direct sum of two toric line bundles.

Example 6. Let $\mathcal{F}$ be a rank 2 toric torsion free sheaf on $\mathbb{P}(2,2,2)$. Again, we will focus on the case $\mathcal{F}$ is a vector bundle. Let $\hat{F}_i$ be the stacky S-families. There are three cases:

(I) Exactly one box summand of $\hat{F}_i$ is nonzero for each $i = 1, 2, 3$. A priori there are $4 \cdot 4 \cdot 4 = 64$ choices of b_i's, but only $4 \cdot 2 \cdot 1 = 8$ of them satisfy the gluing conditions. The rest of this case is exactly as in Example 5.I. In particular, the limiting vector spaces $\mathbb{C}^2$ do not decompose into 1-dimensional spaces with different $\boldsymbol{\mu}_2$-weights.

(II) There is only one $i = 1, 2, 3$ for which only one box summand of $\hat{F}_i$ is nonzero. Say without loss of generality $i = 3$. A priori there are $4 \cdot 6 \cdot 6 = 144$ choices of b_i's, but only 4 of them satisfy the gluing conditions. The rest is as in Example 5.II. In particular, $\mathcal{F}$ is globally the direct sum of two toric line bundles.

(III) For each $i = 1, 2, 3$ two box summands of $\hat{F}_i$ are nonzero. Again, $\mathcal{F}$ is globally the direct sum of two toric line bundles.

Remark 6.2. *Example 6 readily generalizes to a classification of rank 2 toric vector bundles on any* $\mathbb{P}$. *In particular, we can always distinguish types I, II, III. Types II and III are always equivariantly decomposable as a direct sum of two toric line bundles whereas type I could be equivariantly indecomposable. Rank 2 toric torsion free sheaves are obtained from this description by putting 1-dimensional vector spaces* q_i *"in the corners" much like in example 4.*

The reader might have noticed that for all rank 1 and 2 toric torsion free sheaves on $\mathbb{P}$, *we first glued sheaves according to Theorem 5.1 whilst completely ignoring the finite group actions. This is much simpler. Then we noted there exist* unique *finite group actions on the weight spaces which glue this into a toric sheaf. We expect this to be true for toric torsion free sheaves of any rank* $r > 0$. *However, this is not true for general toric coherent sheaves. For example, take* $\mathbb{P}(1,1,2)$ *and take the zero sheaf on charts* $\mathfrak{U}_1, \mathfrak{U}_2$ *and put a single* $\mathbb{C}$ *on an arbitrary lattice point of* $X(\mathbf{T})$ *associated to* $\mathfrak{U}_3$ *and zeros elsewhere. In this case there are no gluing conditions and we can endow* $\mathbb{C}$ *with either weight* 0 *or* 1 *with respect to the* $\boldsymbol{\mu}_2$*-action.*

7. Rank 1 and 2 torsion free sheaves on weighted projective planes

Let $\mathfrak{X}$ be a smooth proper DM stack. Let $\pi : \mathfrak{X} \to X$ be the map to the coarse moduli scheme and assume X is projective. Fix a polarization $\mathcal{O}_X(1)$ on X and a *generating sheaf* $\mathcal{E}$ on $\mathfrak{X}$. A generating sheaf on $\mathfrak{X}$ is a π-very ample locally free sheaf on $\mathfrak{X}$. This notion was introduced by [OS] (see also [Nir, Sect. 2]). In [Nir], Nironi constructs the moduli space M of Gieseker stable (w.r.t. $\mathcal{O}_X(1)$ and $\mathcal{E}$) sheaves on $\mathfrak{X}$ with fixed *modified* Hilbert polynomial. Here the modified Hilbert polynomial of a coherent sheaf $\mathcal{F}$ on $\mathfrak{X}$ is the polynomial

$$P_{\mathcal{O}_X(1),\mathcal{E}}(\mathcal{F}, t) = \chi(\mathfrak{X}, \pi^*\mathcal{O}_X(t) \otimes \mathcal{E}^* \otimes \mathcal{F}).$$

The modified Hilbert polynomial is used to define Gieseker stability in the usual way. The reason $\mathcal{E}$ is introduced is because otherwise $\chi(\mathfrak{X}, \pi^*\mathcal{O}_X(t) \otimes \mathcal{F}) = \chi(X, \mathcal{O}_X(t) \otimes \pi_*\mathcal{F})$ only "sees" the coarse moduli scheme X. The moduli space M is in fact constructed using GIT and it is a quasi-projective $\mathbb{C}$-scheme [Nir, Sect. 6]. In the case there are no strictly Gieseker semistable sheaves M is projective.

Let $\mathbb{P} := \mathbb{P}(a, b, c)$ as before, let $\mathcal{O}_{\mathbf{P}}(1)$ be any polarization on the coarse moduli scheme $\mathbf{P}$ and let $\mathcal{E}$ be any generating sheaf on $\mathbb{P}$. For any class $\mathsf{c} \in K(\mathbb{P})_{\mathbb{Q}}$ of a torsion free sheaf on $\mathbb{P}$, define $M_{\mathcal{E}}(\mathsf{c})$ to be the moduli scheme of μ-stable sheaves on $\mathbb{P}$ of class c. Note that there are two differences with Nironi [Nir]. Firstly, we fix K-group class instead of modified Hilbert polynomial. This is a more refined topological invariant (see also the discussion in the introduction of [Nir]). Secondly, we use μ- instead of Gieseker stability. Here we mean μ-stability defined using the modified Hilbert polynomial, i.e. the linear term of the modified Hilbert polynomial divided by the quadratic term of the modified Hilbert polynomial. The moduli scheme $M_{\mathcal{E}}(\mathsf{c})$ does not depend on the choice of $\mathcal{O}_{\mathbf{P}}(1)$ but does (in general) depend on the choice of generating sheaf $\mathcal{E}$. (For polarizations on $\mathbf{P}$ and a general survey on weighted projective spaces viewed as toric varieties, see e.g. [RT].)

Note that we consider moduli spaces of μ-stable sheaves *only* and we do not consider strictly μ-semistable sheaves. For K-group classes c for which the associated modified Hilbert polynomial satisfies certain numerical constraints on the coefficients, one can ensure there are no strictly μ-semistables around. E.g. see examples 1–4 at the end of section 7.2.

Convention. Throughout this section, we fix the standard polarization $\mathcal{O}_{\mathbf{P}}(1)$ on $\mathbf{P}$ and we choose a generating sheaf $\mathcal{E} := \bigoplus_{u=0}^{E-1} \mathcal{O}_{\mathbb{P}}(-u)$, where E is any positive integer such that the least common multiple m of a, b, c divides E.

Note that this $\mathcal{E}$ admits equivariant structures. This is not the most general choice of generating sheaf on $\mathbb{P}$, but we are getting the nicest formulae in this case. Most of the time, we suppress the dependence on $\mathcal{O}_{\mathbf{P}}(1)$ and $\mathcal{E}$.

7.1. **Rank 1.** In this section, c denotes the class of a rank 1 torsion free sheaf on $\mathbb{P}$ and we denote the corresponding moduli space by $M(\mathsf{c})$. In the rank 1 case $M(\mathsf{c})$ is independent of the choice of polarization as well as generating sheaf because μ-stability is automatic. We are interested in computing the topological Euler characteristics $e(M(\mathsf{c}))$ of these moduli spaces, where we let $\mathsf{c} \in K(\mathbb{P})_{\mathbb{Q}}$ run over all classes of rank 1 torsion free sheaves on $\mathbb{P}$ with fixed first Chern class $\beta \in A^1(\mathbb{P})$

$$\mathsf{G}_\beta(q) = \sum_{c_1(\mathsf{c})=\beta} e(M(\mathsf{c}))q^{\mathsf{c}}.$$

Here q is a formal variable keeping track of the class c.

In practice, q stands for the following variables. We introduce the formal variables $p_0, \ldots, p_{a-1}$, $q_0, \ldots, q_{b-1}$ and $r_0, \ldots, r_{c-1}$ respectively keeping tracks of the classes

$$[\mathcal{O}_{P_1}], \ldots, [\mathcal{O}_{P_1}]g^{a-1}, [\mathcal{O}_{P_2}], \ldots, [\mathcal{O}_{P_2}]g^{b-1} \text{ and } [\mathcal{O}_{P_3}], \ldots, [\mathcal{O}_{P_3}]g^{c-1}$$

in $K(\mathbb{P})_{\mathbb{Q}}$ defined in Section 4.2. In the light of (4), we impose the following relations among these variables:

$$\begin{aligned} p_0 p_d \cdots p_{a-d} &= q_0 q_d \cdots q_{b-d} = r_0 r_d \cdots r_{c-d}, \\ p_1 p_{d+1} \cdots p_{a-d+1} &= q_1 q_{d+1} \cdots q_{b-d+1} = r_1 r_{d+1} \cdots r_{c-d+1}, \\ &\cdots \\ p_{d-1} p_{2d-1} \cdots p_{a-1} &= q_{d-1} q_{2d-1} \cdots q_{b-1} = r_{d-1} r_{2d-1} \cdots r_{c-1}, \\ p_0 p_{d_{12}} \cdots p_{a-d_{12}} &= q_0 q_{d_{12}} \cdots q_{b-d_{12}}, \ldots \\ p_0 p_{d_{13}} \cdots p_{a-d_{23}} &= r_0 r_{d_{13}} \cdots r_{c-d_{13}}, \ldots \\ q_0 q_{d_{23}} \cdots q_{b-d_{23}} &= r_0 r_{d_{23}} \cdots r_{c-d_{23}}, \ldots \end{aligned} \tag{6}$$

As a toric sheaf, $\mathcal{O}_{P_1}$ is described by stacky S-families $\hat{F}_1$, $\hat{F}_2$, $\hat{F}_3$, where $\hat{F}_2 = \hat{F}_3 = 0$ and $\hat{F}_1$ consists of a single vector space $\mathbb{C}$ with trivial $\boldsymbol{\mu}_a$-action at $(0,0) \in X(\mathbf{T}) = \mathbb{Z}^2$ and zeros everywhere else. The toric sheaves $\mathcal{O}_{P_2}$, $\mathcal{O}_{P_3}$ have similar descriptions replacing 1 by 2 and 1 by 3 respectively. Denote by $[\mathcal{O}_{P_1} \otimes \mu]$, $[\mathcal{O}_{P_2} \otimes \nu]$, $[\mathcal{O}_{P_3} \otimes \xi]$ the generators of the Grothendieck groups of the residue gerbes at the points P_1, P_2 and P_3, i.e. $K(B\boldsymbol{\mu}_{\hat{i}}) \cong \mathrm{Rep}(\boldsymbol{\mu}_{\hat{i}})$ for $i = 1, 2, 3$. We use the same notation for the push-forwards of these classes to $K(\mathbb{P})_{\mathbb{Q}}$.

Proposition 7.1. *The K-group classes $[\mathcal{O}_{P_1} \otimes \mu^i]$, $[\mathcal{O}_{P_2} \otimes \nu^j]$, $[\mathcal{O}_{P_3} \otimes \xi^k]$ are equal to respectively $(1-g^b)(1-g^c)g^i$, $(1-g^a)(1-g^c)g^j$, $(1-g^a)(1-g^b)g^k$.*

Proof. We compute $[\mathcal{O}_{P_1} \otimes \mu^i]$ and leave the other cases to the reader. Recall the characterization of toric line bundles $L_{(A,B,C)}$, $A, B, C \in \mathbb{Z}$ of Section 6. Note that $L_{(1,0,0)}$, $L_{(0,1,0)}$, $L_{(0,0,1)}$ generate the equivariant Picard group (Proposition 6.1) and their K-group classes are $g = [\mathcal{O}_{\mathbb{P}}(-1)]$. Using the description of toric line bundles and $\mathcal{O}_{P_1} \otimes \mu^i$ in terms of stacky S-families (Section 6), one readily constructs a $\mathbf{T}$-equivariant short exact sequence

$$0 \longrightarrow L_{(b,c,0)} \longrightarrow L_{(b,0,0)} \oplus L_{(0,c,0)} \longrightarrow L_{(0,0,0)} \longrightarrow \mathcal{O}_{P_1} \longrightarrow 0,$$

where the first map is $v \mapsto (v, v)$ and the second map is $(v, w) \mapsto v - w$. Since $\mathcal{O}_{P_1} \otimes \mu^i \cong \mathcal{O}_{P_1} \otimes L_{(0,0,i)}$ (Section 6), we obtain

$$[\mathcal{O}_{P_1} \otimes \mu^i] = L_{(0,0,i)} - L_{(b,0,i)} - L_{(0,c,i)} + L_{(b,c,i)} = g^i - g^{b+i} - g^{c+i} + g^{b+c+i}.$$

□

Slightly more generally than Proposition 7.1, one can consider a stacky S-family with $\hat{F}_2 = \hat{F}_3 = 0$ and $\hat{F}_1$ of the following form: put a single vector space $\mathbb{C}$ with representation μ^α on the lattice point $i/b(b,0) + j/c(0,c) + I(b,0) + J(0,c) = (i + Ib, j + Jc) \in X(\mathbf{T}) = \mathbb{Z}^2$. The corresponding toric sheaf is easily seen to be

$$\mathcal{O}_{P_1} \otimes L_{(i+Ib, j+Jc, \alpha-i-Ib-j-Jc)},$$

so its K-group class is $(1-g^b)(1-g^c)g^\alpha$. Similarly on the other charts $\mathfrak{U}_2$, $\mathfrak{U}_3$.

Let $\mathcal{F}$ be any rank 1 toric torsion free sheaf on $\mathbb{P}$. The reflexive hull of $\mathcal{F}$ is a toric line bundle $L_{(A,B,C)}$. This is clear from the toric picture of Section 6. The cokernel

$$0 \longrightarrow \mathcal{F} \longrightarrow L_{(A,B,C)} \longrightarrow \mathcal{Q} \longrightarrow 0$$

is a 0-dimensional toric sheaf which can be described by three colored 2D partitions as follows. In each chart $\mathfrak{U}_i$, the stacky S-family of $\mathcal{Q}$ has only one nonzero component $\mathcal{Q}_i$ in its box decomposition. The set

$$\{(\ell_1, \ell_2) \in \mathbb{Z}^2 \mid Q_i(\ell_1, \ell_2) \neq 0\}$$

defines a 2D partition. Suppose $i = 1$ and write uniquely $A = i + Ib$, $B = j + Jc$, $C = k + Ka$ as in Section 6. The 2D partition associated to $\mathcal{Q}_1$ is nonempty if and only if $Q_1(I, J) \neq 0$ and $Q_1(I, J)$ has $\boldsymbol{\mu}_a$-weight

$$A + B + C \bmod a.$$

Since any nonzero element of $Q_1(I, J)$ generates the whole module $\mathcal{Q}|_{\mathfrak{U}_1}$, we know the $\boldsymbol{\mu}_a$-representation of any nonzero $Q_i(\ell_1, \ell_2)$, i.e.

$$A + B + C + \ell_1 b + \ell_2 c \bmod a.$$

This describes a $\mathbb{Z}_a$-coloured 2D partition $\lambda_1(\mathcal{F})$. Similarly, colored partitions $\lambda_2(\mathcal{F})$, $\lambda_3(\mathcal{F})$ arise from the charts $\mathfrak{U}_2$, $\mathfrak{U}_3$. For $i = 1, 2, 3$, we denote by $\Pi_{(A,B,C)}(i)$ the collection of all 2D partitions with $\mathbb{Z}_{\hat{i}}$-colouring arising from some $\lambda_i(\mathcal{F})$, where $\mathcal{F}$ is a rank 1 toric torsion free sheaf on $\mathbb{P}$ with reflexive hull $L_{(A,B,C)}$. Depending on the choice of a, b, c, not all $\hat{i}$ colors might appear.

Example 1. Assume $(a, b, c) = (1, 1, 2)$ and let $A, B, C \in \mathbb{Z}$. The charts $\mathfrak{U}_1$ and $\mathfrak{U}_2$ give rise to the collections $\Pi_{(A,B,C)}(1)$, $\Pi_{(A,B,C)}(2)$ of (uncolored) 2D partitions. Moreover, chart $\mathfrak{U}_3$ gives $\Pi_{(A,B,C)}(3)$, i.e. the collection of $\mathbb{Z}_2$-colored 2D partitions where the first block is 0 if $A+B+C$ is even and 1 if $A+B+C$ is odd:

$(a, b, c) = (1, 1, 2)$

$A + B + C$ even



$(a, b, c) = (1, 1, 2)$

$A + B + C$ odd



Example 2. Assume $(a, b, c) = (2, 2, 2)$ and let $A, B, C \in \mathbb{Z}$. Then $\Pi_{(A,B,C)}(1)$, $\Pi_{(A,B,C)}(2)$, $\Pi_{(A,B,C)}(3)$ are collections of $\mathbb{Z}_2$-colored partitions, but only one color actually appears. This color is 0 when $A + B + C$ is even and 1 when $A + B + C$ is odd

$(a, b, c) = (2, 2, 2)$

$A + B + C$ even



$(a, b, c) = (2, 2, 2)$

$A + B + C$ odd



Back to the general situation. For $i = 1, 2, 3$, the 2D partitions of $\Pi_{(A,B,C)}(i)$ are colored by $\mathbb{Z}_{\hat{i}}$, though not all colors actually have to appear (Example 2). For any $l \in \mathbb{Z}_{\hat{i}}$ and $\lambda \in \Pi_{(A,B,C)}(i)$, we denote by $\#_l\lambda$ the number of boxes of λ with color l. The previous discussion together with Propositions 6.1, 7.1 gives:

Proposition 7.2. *Let $\mathcal{F}$ be a rank 1 toric torsion free sheaf on $\mathbb{P}$. Let $L_{(A,B,C)}$ be the reflexive hull of $\mathcal{F}$ and let $\mathcal{Q}$ be the cokernel of the inclusion $\mathcal{F} \subset L_{(A,B,C)}$. Then $\mathcal{Q}$ is described by three colored partitions $\lambda_i(\mathcal{F}) \in \Pi_{(A,B,C)}(i)$, $i = 1, 2, 3$*

and the K-group class of $\mathcal{F}$ is

$$\left(1-\sum_{i=1}^{3}\sum_{l=0}^{\hat{i}-1}\#_l\lambda_i(\mathcal{F})g^l(1-g^a)(1-g^b)(1-g^c)/(1-g^{\hat{i}})\right)g^{A+B+C}.$$

Suppose $\mathcal{F}$ is a rank 1 toric torsion free sheaf on $\mathbb{P}$ with reflexive hull $L_{(A,B,C)}$. When counting $\mathbf{T}$-fixed points in the proof of the theorem below, we are only interested in the coherent sheaf $\mathcal{F}$ and not its equivariant structure. A toric line bundle $L_{(A',B',C')}$ has trivial underlying line bundle if and only if $A'+B'+C'=0$. By tensoring $\mathcal{F}$ with $L_{(-A,-B,A+B)}$, we change its equivariant structure, but not the underlying coherent sheaf. Therefore, it suffices to only consider rank 1 toric torsion free sheaves $\mathcal{F}$ on $\mathbb{P}$ with reflexive hulls of the form $L_{(0,0,C)}$. In this case, $c_1(\mathcal{F})=-C$. For any $\beta\in A^1(\mathbb{P})=\mathbb{Z}$ and $i=1,2,3$, we define $\Pi_\beta(i):=\Pi_{(0,0,-\beta)}(i)$. The main result of this section is the following theorem.

Theorem 7.3. *For any $\beta\in A^1(\mathbb{P})$, we have*

$$\mathsf{G}_\beta=\left(\sum_{\lambda\in\Pi_\beta(1)}\prod_{l=0}^{a-1}p_l^{\#_l\lambda}\right)\left(\sum_{\lambda\in\Pi_\beta(2)}\prod_{l=0}^{b-1}q_l^{\#_l\lambda}\right)\left(\sum_{\lambda\in\Pi_\beta(3)}\prod_{l=0}^{c-1}r_l^{\#_l\lambda}\right),$$

where the variable p_l,q_l,r_l satisfy relations (6).

Proof. Fix a class $\mathsf{c}\in K(\mathbb{P})_{\mathbb{Q}}$ with $c_1(\mathsf{c})=\beta$. The moduli scheme $M(\mathsf{c})$ has a torus action, which at the level of closed points is given by[14]

$$t\cdot[\mathcal{F}]:=[t^*\mathcal{F}],\ t\in\mathbf{T},\ \mathcal{F}\in M(\mathsf{c}).$$

By localization $e(M(\mathsf{c}))=e(M(\mathsf{c})^{\mathbf{T}})$. The closed points of $M(\mathsf{c})^{\mathbf{T}}$ are exactly the isomorphism classes of rank 1 torsion free sheaves with class c satisfying $t^*\mathcal{F}\cong\mathcal{F}$ for all $t\in\mathbf{T}$. Since any such $\mathcal{F}$ is simple, it admits a $\mathbf{T}$-equivariant structure [Koo1, Prop. 4.4] and this $\mathbf{T}$-equivariant structure is unique up to tensoring by a character of $\mathbf{T}$ [Koo1, Prop. 4.5]. Consequently, $M(\mathsf{c})^{\mathbf{T}}$ is equal to the set[15] of *equivariant* isomorphism classes of rank 1 toric torsion free sheaves with class c, where two such classes $[\mathcal{F}],[\mathcal{F}']$ are identified whenever $\mathcal{F}$ and $\mathcal{F}'$ differ by a character, i.e. whenever $\mathcal{F}\cong\mathcal{F}'\otimes L_{(A,B,C)}$ for some $A+B+C=0$.

Hence any element of $M(\mathsf{c})^{\mathbf{T}}$ can be *uniquely* represented by stacky S-families $\{\hat{F}_i\}_{i=1,2,3}$ of a rank 1 toric torsion free sheaf with reflexive hull $L_{(0,0,-\beta)}$.

---

[14] To construct the action of $\mathbf{T}$ on $M(\mathsf{c})$ as a morphism, one extends the argument of [Koo1, Prop. 4.2] to Nironi's setting [Nir].

[15] Note that this only describes a bijection of finite sets. We are actually dealing with an isomorphism between two finite collections of reduced points by [Koo1, Thm. 4.9] (extended to this setting).

By Proposition 7.2, the elements of $M(\mathsf{c})^{\mathbf{T}}$ are in bijective correspondence to triples of partitions $(\lambda_1, \lambda_2, \lambda_3) \in \Pi_\beta(1) \times \Pi_\beta(2) \times \Pi_\beta(3)$ satisfying

$$\left(1 - \sum_{i=1}^{3} \sum_{l=0}^{\hat{i}-1} \#_l \lambda_i g^l (1-g^a)(1-g^b)(1-g^c)/(1-g^{\hat{i}})\right) g^{-\beta} = \mathsf{c}.$$

□

By Theorem 7.3, the problem of computing Euler characteristics of moduli spaces of the rank 1 torsion free sheaves on $\mathbb{P}$ is reduced to counting colored 2D partitions. Counting colored 2D partitions is a delicate topic. To the authors knowledge, no closed formula exists for the general case. When the action of the cyclic group $\boldsymbol{\mu}_k$ on $\mathbb{C}^2$ is *balanced*, i.e. is of the form $\mu \cdot (x, y) = (\mu x, \mu^{k-1} y)$, there is an elegant formula appearing in the physics literature [DS]. The formula in this case is[16]

$$\sum_{\text{coloured partitions } \lambda} q_0^{\#_0 \lambda} \cdots q_{k-1}^{\#_{k-1}\lambda} = \tag{7}$$

$$\prod_{j>0} \frac{1}{(1-(q_0 \cdots q_{k-1})^j)^k} \sum_{n_1, \ldots, n_{k-1} \in \mathbb{Z}} (q_0 \cdots q_{k-1})^{\sum_i n_i^2 - n_i n_{i+1}} \prod_{r=1}^{k-1} q_{k-r}^{r^2/2 + n_1 r - r/2}.$$

One can count colored 2D partitions keeping track of the number of boxes with color 0 only by setting $q_0 = q$ and $q_1 = \cdots = q_{k-1} = 1$. Then formula (7) is related to the character formula of the affine Lie algebra $\widehat{su}(k)$ (Equation (4.5) in [DS])

$$\sum_{\text{colored partitions } \lambda} q^{\#_0 \lambda} = \frac{q^{k/24}}{\eta(q)} \chi^{\widehat{su}(k)}(0), \tag{8}$$

where $\eta(q)$ is the Dedekind eta function.

Example 1. We only have balanced actions on all three charts of $\mathbb{P}$ for $(a, b, c) = (1, 1, 1)$, $(1, 1, 2)$, and $(1, 2, 3)$ (and permutations). From Theorem 7.3 and formulae (7), (8), we get closed expressions for the generating function as follows.

i) Let $(a, b, c) = (1, 1, 1)$. In this case $p_0 = q_0 = r_0$ by (6) and

$$\mathsf{G}_0 = \frac{1}{\prod_{k>0}(1 - r_0^k)^3}.$$

---

[16] Assuming all colored partitions have color 0 at the origin.

ii) Let $(a,b,c)=(1,1,2)$. In this case we have $p_0=q_0=r_0r_1$ by (6) and
$$\mathsf{G}_0=\frac{1}{\prod_{k>0}(1-(r_0r_1)^k)^4}\sum_{k\in\mathbb{Z}} r_0^{k^2}r_1^{k^2+k}.$$
Putting $r_1=1$ and $r_0=q$ in G_0, one gets
$$\mathsf{G}_0(q)=\frac{q^{1/6}}{\eta(q)^4}\chi^{\widehat{su}(2)}(0)=\frac{q^{1/6}}{\eta(q)^4}\theta_3(q),$$
where $\theta_3(q)$ is a Jacobi theta function. Geometrically, the power of q keeps track of the Euler characteristic $\chi(\mathcal{F}^{**}/\mathcal{F})$ of our rank 1 torsion free sheaves $\mathcal{F}$, where $\mathcal{F}^{**}$ is the reflexive hull of $\mathcal{F}$. This can be seen by noting that $\chi(\mathcal{O}_{P_1}\otimes\mu^i)=\delta_{0i}$, $\chi(\mathcal{O}_{P_2}\otimes\nu^j)=\delta_{0j}$ and $\chi(\mathcal{O}_{P_3}\otimes\xi^k)=\delta_{0k}$.

iii) Let $(a,b,c)=(1,2,3)$. In this case $p_0=q_0q_1=r_0r_1r_2$ by (6) and
$$\mathsf{G}_0=\frac{1}{\prod_{k>0}(1-(r_0r_1r_2)^k)^6}\sum_{k\in\mathbb{Z}}(r_0r_1r_2)^{k^2}q_1^k\sum_{k,l\in\mathbb{Z}}r_0^{k^2-kl+l^2}r_1^{k^2+2k+1-kl+l^2}r_2^{k^2+k-kl+l^2}.$$
Setting $q_1=r_1=r_2=1$ and $p_0=q_0=r_0=q$ in G_0 as in the previous case, one gets
$$\mathsf{G}_0(q)=\frac{q^{1/4}}{\eta(q)^6}\chi^{\widehat{su}(2)}(0)\chi^{\widehat{su}(3)}(0)=\frac{q^{1/4}}{\eta(q)^6}\theta_3(q)(\theta_3(q)\theta_3(q^3)+\theta_2(q)\theta_2(q^3)),$$
where $\theta_2(q)$, $\theta_3(q)$ are Jacobi theta functions.

Example 2. Let $(a,b,c)=(1,c,c)$ with $c\geq 2$. In this case there is also a nice formula. Relations (6) give $p_0=q_0\cdots q_{c-1}$ and $q_i=r_i$. The following formula can be easily proved
$$\mathsf{G}_0=\frac{1}{\prod_{k>0}(1-(r_0\cdots r_{c-1})^k)}\frac{1}{(\prod_{k>0}\prod_{i=0}^{c-2}(1-r_0\cdots r_i(r_0\cdots r_{c-1})^{k-1}))^2}.$$

Example 3. For $\mathbb{P}(1,1,3)$, we encounter a non-balanced $\boldsymbol{\mu}_3$-action in the chart $\mathfrak{U}_3$. We do not know a closed formula for G_0 in this example. Relations (6) give $p_0=q_0=r_0r_1r_2$. The first few terms of the expansion are
$$\mathsf{G}_0=\frac{(1+r_0+2r_0r_1+2r_0r_1r_2+r_0r_1^2+2r_0^2r_1r_2+3r_0r_1^2r_3+O(5))}{\prod_{k>0}(1-(r_0r_1r_2)^k)^2}.$$

7.2. **Rank 2.** In this section, $\mathsf{c}\in K(\mathbb{P})_{\mathbb{Q}}$ denotes the class of a rank 2 torsion free sheaf on $\mathbb{P}:=\mathbb{P}(a,b,c)$. We fix $\mathcal{O}_{\mathbf{P}}(1)$ and $\mathcal{E}:=\bigoplus_{u=0}^{E-1}\mathcal{O}_{\mathbb{P}}(-u)$ as in the beginning of this section. The moduli scheme of μ-stable sheaves on $\mathbb{P}$ with class c is denoted by $M_{\mathcal{E}}(\mathsf{c})$ (see the beginning of this section). Unlike the rank 1 case, not all rank 2 torsion free sheaves on $\mathbb{P}$ are μ-stable. We denote by $N_{\mathcal{E}}(\mathsf{c})\subset M_{\mathcal{E}}(\mathsf{c})$ the open subset of the μ-stable vector bundles.

Denote the inertia stack by $I\mathbb{P}$. Using our explicit calculations together with the results of [BCS, BH], one can see that the Chern character map (see Section 4.2)

$$\widetilde{\mathrm{ch}} : K(\mathbb{P})_{\mathbb{Q}} \longrightarrow A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty},$$

is a ring isomorphism. Therefore, fixing a class in $\mathsf{c} \in K(\mathbb{P})_{\mathbb{Q}}$ is equivalent to fixing an element $\widetilde{\mathrm{ch}} \in A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$. Recall that the set F indexes the components of the inertia stack $I\mathbb{P}$ (see Section 4.2). For a fixed $f \in F$ corresponding to component Z, let $\widetilde{\mathrm{ch}}_f$ denote the part of $\widetilde{\mathrm{ch}}$ taking values in $A^*(Z)_{\boldsymbol{\mu}_\infty}$. For notational convenience, we define $(\widetilde{\mathrm{ch}}_f)^k := (\widetilde{\mathrm{ch}}_f)_{\dim Z-k} \in A^{\dim Z-k}(Z)_{\boldsymbol{\mu}_\infty}$ to be its dimension k part. In this context, we refer to k as the codegree. Fix $\alpha \in A^0(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ and $\beta \in A^1(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ (parts of) the Chern character of a rank 2 torsion free sheaf on $\mathbb{P}$. Define the generating functions

$$\mathsf{H}_{\alpha,\beta}(q) := \sum_{\substack{\widetilde{\mathrm{ch}}^2(\mathsf{c}) = \alpha \\ \widetilde{\mathrm{ch}}^1(\mathsf{c}) = \beta}} e(M_{\mathcal{E}}(\mathsf{c}))q^{\mathsf{c}}, \qquad \mathsf{H}^{vb}_{\alpha,\beta}(q) := \sum_{\substack{\widetilde{\mathrm{ch}}^2(\mathsf{c}) = \alpha \\ \widetilde{\mathrm{ch}}^1(\mathsf{c}) = \beta}} e(N_{\mathcal{E}}(\mathsf{c}))q^{\mathsf{c}}.$$

So in terms of $\widetilde{\mathrm{ch}}$, these generating functions sum over all 0-dimensional (or codegree 0) parts $(\widetilde{\mathrm{ch}}_f)^0$.

Our goal is to determine the generating function for the Euler characteristics $M_{\mathcal{E}}(\mathsf{c})$. By the following lemma, this can be achieved by determining the generating functions for the Euler characteristics of $N_{\mathcal{E}}(\mathsf{c})$ and of the moduli spaces of rank 1 torsion free sheaves on $\mathfrak{U}_1, \mathfrak{U}_2, \mathfrak{U}_3$. The latter generating functions, which we denote by $\mathsf{G}_{\mathfrak{U}_i}(q)$, are known by the result of Section 7.1. The lemma is a variation on a result by Göttsche and Yoshioka [Got, Prop. 3.1].

Lemma 7.4. *The following holds:* $\mathsf{H}_{\alpha,\beta}(q) = \mathsf{H}^{vb}_{\alpha,\beta}(q) \prod_{i=1}^3 \mathsf{G}_{\mathfrak{U}_i}(q)^2$.

Proof. Embedding into the reflexive hull of a rank 2 torsion free sheaf $\mathcal{F}$ on $\mathbb{P}$ gives the short exact sequence $0 \to \mathcal{F} \to \mathcal{F}^{**} \to \mathcal{Q} \to 0$, where $\mathcal{Q}$ is a 0-dimensional sheaf on $\mathbb{P}$. Since $\mathbb{P}$ is a smooth DM stack of dimension two $\mathcal{F}^{**}$ is a rank 2 vector bundle.

Let $\mathrm{Quot}_{\mathbb{P}}(\mathcal{V}, \mathsf{c})$ be the projective scheme of quotients of a vector bundle $\mathcal{V}$ on $\mathbb{P}$ representing class $\mathsf{c} \in K(\mathbb{P})_{\mathbb{Q}}$ (see [OS] for the construction of these Quot schemes). Fix $\mathsf{c} \in K(\mathbb{P})_{\mathbb{Q}}$ the class of a rank 2 torsion free sheaf on $\mathbb{P}$. By the previous short exact sequence, at the level of closed points, $M_{\mathcal{E}}(\mathsf{c})$ can be stratified by $\mathrm{Quot}_{\mathbb{P}}(\mathcal{V}, \mathsf{c}')$, $[\mathcal{V}] \in N_{\mathcal{E}}(\mathsf{c}'')$ with $\mathsf{c}' + \mathsf{c}'' = \mathsf{c}$. Hence at the level of closed points, $M_{\mathcal{E}}(\mathsf{c})$ is in bijective correspondence with the disjoint union.

$$\coprod_{\mathsf{c}'+\mathsf{c}''=\mathsf{c}} \ \coprod_{[\mathcal{V}] \in N_{\mathcal{E}}(\mathsf{c}'')} \mathrm{Quot}_{\mathbb{P}}(\mathcal{V}, \mathsf{c}').$$

Let $\mathcal{V}$ be a rank 2 vector bundle on $\mathbb{P}$ with $\mathbf{T}$-equivariant structure. In order to determine $e(\mathrm{Quot}_{\mathbb{P}}(\mathcal{V}, \mathsf{c}'))$ for any 0-dimensional c', we can count the number of $\mathbf{T}$-fixed points using the $\mathbf{T}$-action on $\mathbb{P}$. This $\mathbf{T}$-action lifts to $\mathrm{Quot}_{\mathbb{P}}(\mathcal{V}, \mathsf{c}')$ (see proof of [Koo1, Prop. 4.1]). Since $\mathcal{Q}$ is 0-dimensional with support at $P_1, P_2, P_3 \in \mathbb{P}$, giving a $\mathbf{T}$-fixed quotient $\mathcal{V} \to \mathcal{Q} \to 0$ is equivalent to giving the quotients $\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2} \to \mathcal{Q}|_{\mathfrak{U}_i} \to 0$ for $i = 1, 2, 3$. So at the level of closed points, $\mathrm{Quot}_{\mathbb{P}}^{\mathbf{T}}(\mathcal{V}, \mathsf{c}')$ is in bijective correspondence with the disjoint union

$$\coprod_{\mathsf{c}_1+\mathsf{c}_2+\mathsf{c}_3=\mathsf{c}'} \prod_{i=1}^{3} \mathrm{Quot}_{\mathfrak{U}_i}^{\mathbf{T}}(\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2}, \mathsf{c}_i).$$

We are reduced to determine $e(\mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2}, \mathsf{c}_i))$. We can use the $\mathbb{C}^{*2}$-action on $\mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2}, \mathsf{c}_i)$, which scales the factors of $\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2}$. At the level of closed points, it can be easily seen that the fixed set of this action is in bijective correspondence with the disjoint union

$$\coprod_{\mathsf{c}_i'+\mathsf{c}_i''=\mathsf{c}_i} \mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}, \mathsf{c}_i') \times \mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}, \mathsf{c}_i'').$$

But $\mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}, \mathsf{c}_i'')$ is isomorphic to the moduli space of rank 1 torsion free sheaves on $\mathfrak{U}_i$ with class $-\mathsf{c}_i''$. This proves $\sum_{\mathsf{c}_i} e(\mathrm{Quot}_{\mathfrak{U}_i}(\mathcal{O}_{\mathfrak{U}_i}^{\oplus 2}, \mathsf{c}_i)) q^{\mathsf{c}_i} = \mathsf{G}_{\mathfrak{U}_i}(q)^2$, which finishes the proof of the lemma. $\square$

The rest of the paper is devoted to the study of rank 2 toric vector bundles on $\mathbb{P}$ and the generating function $\mathsf{H}_{\alpha,\beta}^{vb}(q)$. In Section 6, we classified three types of toric vector bundles on $\mathbb{P}$: type I, II and III. Since types II and III are always decomposable, they are never μ-stable. Hence we only need to consider type I rank 2 toric vector bundles $\mathcal{F}$ on $\mathbb{P}$. The stacky S-family of $\mathcal{F}$ is entirely determined by integers $A_1, A_2, A_3 \in \mathbb{Z}$ and $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ satisfying $b \mid \Delta_1$, $c \mid \Delta_2$, $a \mid \Delta_3$ and an element $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$ (see Section 6). We want to express the K-group class of $[\mathcal{F}]$ in terms of this data only. For any two 1-dimensional vector spaces $p, q \in \mathbb{P}^1$, the Kronecker delta δ_{pq} is 1 if $p = q$ and 0 if $p \neq q$.

Proposition 7.5. *Let $\mathcal{F}$ be a type I rank 2 toric vector bundle on $\mathbb{P}$. Let its stacky S-families be described by $A_1, A_2, A_3 \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ satisfying $b \mid \Delta_1$, $c \mid \Delta_2$, $a \mid \Delta_3$ and $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$. Then the K-group class of $\mathcal{F}$ is*

$$\begin{aligned}\big(1 + g^{\Delta_1+\Delta_2+\Delta_3} &- (1-g^{\Delta_1})(1-g^{\Delta_2})(1-\delta_{p_1p_2}) - (1-g^{\Delta_2})(1-g^{\Delta_3})(1-\delta_{p_2p_3}) \\ &- (1-g^{\Delta_3})(1-g^{\Delta_1})(1-\delta_{p_3p_1})\big) g^{A_1+A_2+A_3}.\end{aligned}$$

Proof. We make use of the following observation. Let $\mathcal{G}$ be another toric torsion free sheaf with stacky S-families $\hat{G}_1$, $\hat{G}_2$, $\hat{G}_3$. Assume

$$\dim({}_bG_i(\ell_1,\ell_2)_l) = \dim({}_bF_i(\ell_1,\ell_2)_l), \tag{9}$$

for all $i = 1,2,3$, $b \in \mathcal{B}_{\mathbf{T}}$, $\ell_1, \ell_2 \in \mathbb{Z}$ and $l \in \mathbb{Z}_{\hat{i}}$. Then we claim $[\mathcal{F}] = [\mathcal{G}] \in K(\mathbb{P})_{\mathbb{Q}}$. This is proved in the lemma after this proof using equivariant dévissage.

Assume without lost of generality $A_1 = A_2 = A_3 = 0$. The general case follows from tensoring with $L_{(A_1,A_2,A_3)}$. We construct a toric sheaf $\mathcal{G}$ satisfying Equation (9), but being of a simpler form than $\mathcal{F}$. We take $\mathcal{G}$ to be an equivariant subsheaf of $L_{(0,0,0)} \oplus L_{(\Delta_1,\Delta_2,\Delta_3)}$. In each chart, such a sheaf only has a nonzero box summand over $(0,0)$. For each $i = 1,2,3$, define[17]

$$_{(0,0)}G_i(\ell_1,\ell_2) := {}_{(0,0)}L_{(0,0,0),i}(\ell_1,\ell_2) \oplus {}_{(0,0)}L_{(\Delta_1,\Delta_2,\Delta_3),i}(\ell_1,\ell_2),$$

in the following regions

$$\begin{aligned}&\{(\ell_1,\ell_2) \mid \ell_1 \geq \Delta_i \text{ or } \ell_2 \geq \Delta_{i+1}\},\\ &\{(\ell_1,\ell_2) \mid \ell_1 < \Delta_i \text{ and } \ell_2 < \Delta_{i+1}\}, \text{ if } p_i = p_{i+1}\end{aligned}$$

and ${}_{(0,0)}G_i(\ell_1,\ell_2) = 0$ for all remaining (ℓ_1,ℓ_2). From this description it is easy to see that Equation (9) is satisfied and $[\mathcal{G}]$ is equal to

$$\begin{aligned}1 + g^{\Delta_1+\Delta_2+\Delta_3} - (1-\delta_{p_1p_2}) &\sum_{I=0}^{\Delta_1/b-1} \sum_{J=0}^{\Delta_2/c-1} [\mathcal{O}_{P_1} \otimes \mu^{Ib+Jc}]\\ - (1-\delta_{p_2p_3}) &\sum_{J=0}^{\Delta_2/c-1} \sum_{K=0}^{\Delta_3/a-1} [\mathcal{O}_{P_2} \otimes \nu^{Jc+Ka}]\\ - (1-\delta_{p_3p_1}) &\sum_{K=0}^{\Delta_3/a-1} \sum_{I=0}^{\Delta_1/b-1} [\mathcal{O}_{P_3} \otimes \xi^{Ka+Ib}].\end{aligned}$$

The sums can be computed using Proposition 7.1. What left is the following lemma. □

Lemma 7.6. *Let $\mathcal{F}$, $\mathcal{G}$ be toric torsion free sheaves on $\mathbb{P}$ and assume their stacky S-families only possibly differ in the maps between the fine-graded weight spaces. Then $[\mathcal{F}] = [\mathcal{G}] \in K(\mathbb{P})_{\mathbb{Q}}$.*

Proof. We give the argument for toric torsion free sheaves on $\mathbb{C}^2$ with standard $\mathbf{T}$-action. The reader can easily generalize the argument to $\mathbb{P}$. Take $(1,0)$, $(0,1)$ as a basis for the character group $M = X(\mathbf{T})$. Let $\mathcal{F}$, $\mathcal{G}$ be toric torsion free sheaves on $\mathbb{C}^2$ and denote their S-families by $\hat{F}$, $\hat{G}$. Assume $\mathcal{F}, \mathcal{G}$ only possibly differ in the maps between their weight spaces. We apply *equivariant dévissage*. Since both sheaves are finitely generated, there are integers A, B such that all homogeneous generators of $H^0(\mathcal{F})$ and $H^0(\mathcal{G})$ lie in $(-\infty, A) \times (-\infty, B)$. In particular, for all $\ell_1 \geq A$ and $\ell_2 \geq B$ we have $F_1(\ell_1, \ell_2) = G_1(\ell_1, \ell_2) = \mathbb{C}^2$ and the maps between the weight spaces are the identity. Define the S-family

---

[17] Here $\Delta_4 := \Delta_1$ and $p_4 := p_1$.

$$\begin{aligned} S_1(\ell_1, \ell_2) &:= F(\ell_1, \ell_2), \text{ for } \ell_1 \geq A \text{ and } \ell_2 \geq B, \\ S_1(\ell_1, \ell_2) &:= 0, \text{ otherwise.} \end{aligned}$$

At the level of S-families, $S_1 \subset F$ and we denote the quotient by Q_1. These correspond to a toric sheaf $\mathcal{S}_1 \subset \mathcal{F}$ with quotient sheaf $\mathcal{Q}_1$ and we get $[\mathcal{F}] = [\mathcal{S}_1] + [\mathcal{Q}_1]$. Repeating the same argument for $\mathcal{G}$, we obtain $[\mathcal{G}] = [\tilde{\mathcal{S}}_1] + [\tilde{\mathcal{Q}}_1] = [\mathcal{S}_1] + [\tilde{\mathcal{Q}}_1]$ since $\tilde{\mathcal{S}}_1 = \mathcal{S}_1$. (Aside: $\mathcal{S}_1 \cong \mathcal{O}_{\mathbb{C}^2}^{\oplus r}$, where r is the rank.)

We shift focus to $\mathcal{Q}_1$, $\tilde{\mathcal{Q}}_1$. Define the S-family

$$\begin{aligned} S_2(\ell_1, \ell_2) &:= F(\ell_1, \ell_2), \text{ for } \ell_1 = A-1 \text{ and } \ell_2 \geq B, \\ S_2(\ell_1, \ell_2) &:= 0, \text{ otherwise.} \end{aligned}$$

This gives a toric sheaf $\mathcal{S}_2 \subset \mathcal{Q}_1$ with quotient $\mathcal{Q}_2$ and $[\mathcal{Q}_1] = [\mathcal{S}_2] + [\mathcal{Q}_2]$. Repeating the same argument for $\tilde{\mathcal{Q}}_1$, we obtain $[\tilde{\mathcal{Q}}_1] = [\tilde{\mathcal{S}}_2] + [\tilde{\mathcal{Q}}_2] = [\mathcal{S}_2] + [\tilde{\mathcal{Q}}_2]$ since $\tilde{\mathcal{S}}_2 = \mathcal{S}_2$. (Aside: $\mathcal{S}_2 \cong \mathcal{O}_{\mathbb{C}}^{\oplus s}$ for some s.)

Repeating the argument a finite number of times, we "peel off" the region where $\ell_1 \geq A$ or $\ell_2 \geq B$. What is left are toric sheaves $\mathcal{Q}_n$, $\tilde{\mathcal{Q}}_n$ defined by the S-families

$$\begin{aligned} &Q_n(\ell_1, \ell_2) := F(\ell_1, \ell_2), \ \tilde{Q}_n(\ell_1, \ell_2) := G(\ell_1, \ell_2), \text{ for } \ell_1 \leq A-1 \text{ and } \ell_2 \leq B-1, \\ &Q_n(\ell_1, \ell_2) := 0, \ \tilde{Q}_n(\ell_1, \ell_2) := 0, \text{ otherwise.} \end{aligned}$$

It suffices to show $[\mathcal{Q}_n] = [\tilde{\mathcal{Q}}_n]$.

For this, define the S-family

$$\begin{aligned} S_{n+1}(\ell_1, \ell_2) &:= F(\ell_1, \ell_2), \text{ for } (\ell_1, \ell_2) = (A-1, B-1), \\ S_{n+1}(\ell_1, \ell_2) &:= 0, \text{ otherwise.} \end{aligned}$$

This gives a toric sheaf $\mathcal{S}_{n+1} \subset \mathcal{Q}_n$ with quotient $\mathcal{Q}_{n+1}$ and $[\mathcal{Q}_n] = [\mathcal{S}_{n+1}] + [\mathcal{Q}_{n+1}]$. Repeating the same argument for $\tilde{\mathcal{Q}}_{n+1}$, we obtain $[\tilde{\mathcal{Q}}_n] = [\tilde{\mathcal{S}}_{n+1}] + [\tilde{\mathcal{Q}}_{n+1}] = [\mathcal{S}_{n+1}] + [\tilde{\mathcal{Q}}_{n+1}]$ since $\tilde{\mathcal{S}}_{n+1} = \mathcal{S}_{n+1}$. (Aside: $\mathcal{S}_{n+1} \cong \mathcal{O}_0^{\oplus t}$ for some t.)

Repeating the final part a finite number of times, the lemma is proved. □

We can now classify all rank 2 μ-stable toric vector bundles on $\mathbb{P}$. Recall that we fix the standard polarization $\mathcal{O}_{\mathbf{P}}(1)$ on $\mathbf{P}$ and a generating sheaf of the form $\mathcal{E} := \bigoplus_{u=0}^{E-1} \mathcal{O}_{\mathbb{P}}(-u)$ on $\mathbb{P}$, where $E \in \mathbb{Z}_{>0}$ is divisible by the least common multiple m of a, b, c. Let $P_{\mathcal{E}}(-,t)$ be the modified Hilbert polynomial with respect to the polarization $\mathcal{O}_{\mathbf{P}}(1)$ and generating sheaf $\mathcal{E}$ as fixed at the beginning of this section. Note that $\mathcal{O}_{\mathbf{P}}(1)$ pulls back to $\mathcal{O}_{\mathbb{P}}(m)$ on $\mathbb{P}$. (See Section 4.2 for the definition of the common divisors d, d_{ij} and the common multiple m.) The modified slope μ of any torsion free sheaf on $\mathbb{P}$ is defined to be the ratio of the linear and quadratic terms of its modified Hilbert polynomial.

Proposition 7.7. *Let $\mathcal{F}$ be a type I rank 2 toric vector bundle on $\mathbb{P}$ and let its stacky S-families be described by $A_1, A_2, A_3 \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ satisfying $b \mid \Delta_1$, $c \mid \Delta_2$, $a \mid \Delta_3$ and $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$. Then $\mathcal{F}$ is μ-stable if and only if $\Delta_1, \Delta_2, \Delta_3 > 0$, p_1, p_2, p_3 are mutually distinct and*

$$\Delta_1 < \Delta_2 + \Delta_3, \ \Delta_2 < \Delta_1 + \Delta_3, \ \Delta_3 < \Delta_1 + \Delta_2.$$

Proof. We start by observing that $\mathcal{F}$ is μ-semistable if and only if $\mu(\mathcal{G}) \leq \mu(\mathcal{F})$ for any *toric* subsheaf $\mathcal{G}$ with $0 < \mathrm{rk}(\mathcal{G}) < \mathrm{rk}(\mathcal{F})$. This is immediate by noting that the Harder-Narasimhan filtration of $\mathcal{F}$ is $\mathbf{T}$-equivariant. It is also true that $\mathcal{F}$ is μ-stable if and only if $\mu(\mathcal{G}) < \mu(\mathcal{F})$ for any *toric* subsheaf $\mathcal{G}$ with $0 < \mathrm{rk}(\mathcal{G}) < \mathrm{rk}(\mathcal{F})$. This statement is more subtle and is proved in the case of smooth toric varieties in [Koo1, Prop. 4.13]. The proof extends to our current setting.

Note that $\mathcal{F}$ can only be indecomposable if $\Delta_1, \Delta_2, \Delta_3 > 0$ and p_1, p_2, p_3 are mutually distinct so we assume this from now on. From the toric description of $\mathcal{F}$ in terms of stacky S-families, it is clear that $\mathcal{F}$ has three toric line bundles $\mathcal{L}_1$, $\mathcal{L}_2$, $\mathcal{L}_3$ as subsheaves generated p_1, p_2, p_3. Moreover, any rank 1 toric subsheaf is contained in one of them. Their stacky S-families and K-group classes are easily computed

$$[\mathcal{L}_1] = g^{\Delta_2+\Delta_3} g^{A_1+A_2+A_3}, \ [\mathcal{L}_2] = g^{\Delta_1+\Delta_3} g^{A_1+A_2+A_3}, \ [\mathcal{L}_3] = g^{\Delta_1+\Delta_2} g^{A_1+A_2+A_3}.$$

The K-group class of $\mathcal{F}$ was computed in Proposition 7.5. Since all these K-group classes are expressed as sums of elements g^r, we can use Corollary 4.4 to compute the top two coefficients of the modified Hilbert polynomials and hence the modified slopes

$$\mu_{\mathcal{F}} = \frac{1}{m}(a+b+c-(\Delta_1+\Delta_2+\Delta_3) - 2(A_1+A_2+A_3)) + \frac{2}{em}\sum_{i=1}^{e} u_i,$$

$$\mu_{\mathcal{L}_i} = \frac{1}{m}(a+b+c-2(\Delta_j+\Delta_k+A_1+A_2+A_3)) + \frac{2}{em}\sum_{i=1}^{e} u_i,$$

for any $i = 1, 2, 3$ and distinct $j, k \in \{1,2,3\} - \{i\}$. Since any other rank 1 toric subsheaf is contained in some $\mathcal{L}_i$ and has modified slope $\leq \mu_{\mathcal{L}_i}$, it suffices to test μ-stability for $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$ and the proposition follows. □

From Proposition 7.5, it is easy to compute the Chern character of any type I rank 2 toric vector bundle on $\mathbb{P}$. Since we are only interested in indecomposable vector bundles, we assume all $\Delta_i > 0$ and p_i are mutually distinct. Moreover, since we are only interested in $\mathbf{T}$-equivariant structures up to tensoring by a character, we assume $A_1 = A_2 = 0$. See Section 4.2 for a discussion of the Chern character map $\widetilde{\mathrm{ch}} : K(\mathbb{P})_{\mathbb{Q}} \to A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$.

Lemma 7.8. *Let $\mathcal{F}$ be a type I rank 2 toric vector bundle on $\mathbb{P}$. Let its stacky S-families be described by $A_1, A_2, A_3 \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{\geq 0}$ satisfying $b \mid \Delta_1$, $c \mid \Delta_2$, $a \mid \Delta_3$ and $(p_1, p_2, p_3) \in (\mathbb{P}^1)^3$. Assume all $\Delta_i > 0$, p_i are mutually distinct and $A_1 = A_2 = 0$, $A_3 = A$. Then $\widetilde{\mathrm{ch}}(\mathcal{F})$ is given by*

$$e^{-2\pi\sqrt{-1}fA}\Big\{2-\big(2A+\Delta_1+\Delta_2+\Delta_3\big)x+\big(A^2+(\Delta_1+\Delta_2+\Delta_3)A+\frac{1}{2}(\Delta_1^2+\Delta_2^2+\Delta_3^2)\big)x^2\Big\}$$

for any $f \in D$;

$$e^{-2\pi\sqrt{-1}fA}\Big\{(1+e^{-2\pi\sqrt{-1}f\Delta_j})-\big((1+e^{-2\pi\sqrt{-1}f\Delta_j})A+\Delta_i+e^{-2\pi\sqrt{-1}f\Delta_j}\Delta_j+\Delta_k\big)x\Big\},$$

for any $(i,j) = (1,2), (2,3), (3,1)$, $f \in D_{ij}$ and $k \in \{1,2,3\} \setminus \{i,j\}$;

$$e^{-2\pi\sqrt{-1}fA}\left\{e^{-2\pi\sqrt{-1}f\Delta_i} + e^{-2\pi\sqrt{-1}f\Delta_{i+1}}\right\},$$

for any $i = 1,2,3$ and $f \in D_i$, where $\Delta_4 := \Delta_1$.

Fix $\alpha \in A^0(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ and $\beta \in A^1(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ (parts of) the Chern character of a rank 2 vector bundle on $\mathbb{P}$. Let $\widetilde{\mathrm{ch}}(A, \Delta_1, \Delta_2, \Delta_3) \in A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ be the formula for the Chern character as appearing in the previous lemma. Consider the following system of equations in the variables $A \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{>0}$

$$\widetilde{\mathrm{ch}}^2(A, \Delta_1, \Delta_2, \Delta_3) = \alpha,$$
$$\widetilde{\mathrm{ch}}^1(A, \Delta_1, \Delta_2, \Delta_3) = \beta.$$

Note that the $\widetilde{\mathrm{ch}}^2$ equation does not depend on $\Delta_1, \Delta_2, \Delta_3$. Moreover, the $\widetilde{\mathrm{ch}}^2$ equation is automatically satisfied in the case of a trivial gerbe structure, i.e. when $d = 1$. In the case of a non-trivial gerbe structure, the $\widetilde{\mathrm{ch}}^2$ equation is equivalent to fixing $A \equiv \lambda \bmod d$ for some $\lambda \in \{0, \ldots, d-1\}$. Next, look at the $f = 0 \in D$ part of the $\widetilde{\mathrm{ch}}^1$ equation and let β_0 be the $f = 0 \in D$ part of β. This equation only has a solution if $2 \mid \beta_0 + \Delta_1 + \Delta_2 + \Delta_3$, in which case

$$A = -\frac{1}{2}\left(\beta_0 + \Delta_1 + \Delta_2 + \Delta_3\right).$$

What remains are the $\widetilde{\mathrm{ch}}^1$ equations for $f \in D \setminus 0$, which we analyze further in examples.

For each $f \in D$ we introduce a formal variable p_f. Likewise, for each $i < j \in \{1,2,3\}$ and $f \in D_{ij}$ we introduce a formal variable $q_{ij,f}$. Finally, for each $i \in \{1,2,3\}$ and $f \in D_i$ we introduce a formal variable $r_{i,f}$. Note that by working with Chern character instead of K-group class, we know these variables are independent. We can now explicitly compute the generating function $\mathsf{H}^{vb}_{\alpha,\beta}$.

Theorem 7.9. *For any $\alpha \in A^0(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ and $\beta \in A^1(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ (part of) the Chern character of a rank 2 vector bundle on $\mathbb{P}$, the generating function $\mathsf{H}^{vb}_{\alpha,\beta}$ is given by*

$$\sum_{\substack{\Delta_1,\Delta_2,\Delta_3 \in \mathbb{Z}_{>0},\, A \in \mathbb{Z} \\ b \mid \Delta_1,\, c \mid \Delta_2,\, a \mid \Delta_3 \\ \widetilde{\mathrm{ch}}^2(A,\Delta_1,\Delta_2,\Delta_3) = \alpha \\ \widetilde{\mathrm{ch}}^1(A,\Delta_1,\Delta_2,\Delta_3) = \beta \\ \Delta_1 < \Delta_2 + \Delta_3 \\ \Delta_2 < \Delta_1 + \Delta_3 \\ \Delta_3 < \Delta_1 + \Delta_2}} \prod_{f \in D} p_f^{\widetilde{\mathrm{ch}}^0(A,\Delta_1,\Delta_2,\Delta_3)_f} \prod_{\substack{i < j \in \{1,2,3\} \\ f \in D_{ij}}} q_{ij,f}^{\widetilde{\mathrm{ch}}^0(A,\Delta_1,\Delta_2,\Delta_3)_f} \prod_{\substack{i \in \{1,2,3\} \\ f \in D_i}} r_{i,f}^{\widetilde{\mathrm{ch}}^0(A,\Delta_1,\Delta_2,\Delta_3)_f},$$

where $\widetilde{\mathrm{ch}}^0(A,\Delta_1,\Delta_2,\Delta_3)$ is the explicit formula for the codegree 0 part of the Chern character appearing in Lemma 7.8 and $\widetilde{\mathrm{ch}}^0(A,\Delta_1,\Delta_2,\Delta_3)_f$, $f \in F$ are its various components.

Proof. We start as in the rank 1 case. Fix a Chern character $\widetilde{\mathrm{ch}} \in A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ of a rank 2 vector bundle on $\mathbb{P}$. The moduli scheme $N(\widetilde{\mathrm{ch}})$ of μ-stable vector bundles on $\mathbb{P}$ with Chern character $\widetilde{\mathrm{ch}}$ has a torus action, which at the level of closed points is given by

$$t \cdot [\mathcal{F}] := [t^*\mathcal{F}],\ t \in \mathbf{T},\ \mathcal{F} \in N(\widetilde{\mathrm{ch}}).$$

By localization, $e(N(\widetilde{\mathrm{ch}})) = e(N(\widetilde{\mathrm{ch}})^{\mathbf{T}})$. As a set, $N(\widetilde{\mathrm{ch}})^{\mathbf{T}}$ is the collection of isomorphism classes of rank 2 μ-stable vector bundles $\mathcal{F}$ on $\mathbb{P}$ with Chern character $\widetilde{\mathrm{ch}}$ satisfying $t^*\mathcal{F} \cong \mathcal{F}$ for all $t \in \mathbf{T}$. Since any such $\mathcal{F}$ is simple, it admits a $\mathbf{T}$-equivariant structure [Koo1, Prop. 4.4] and this $\mathbf{T}$-equivariant structure is unique up to tensoring by a character of $\mathbf{T}$ [Koo1, Prop. 4.5]. Consequently, as a set, $N(\widetilde{\mathrm{ch}})^{\mathbf{T}}$ is the collection of *equivariant* isomorphism classes of μ-stable rank 2 toric vector bundles on $\mathbb{P}$ with class $\widetilde{\mathrm{ch}}$, where two such classes $[\mathcal{F}], [\mathcal{F}']$ are identified whenever $\mathcal{F}$ and $\mathcal{F}'$ differ by a character, i.e. whenever $\mathcal{F} \cong \mathcal{F}' \otimes L_{(A,B,C)}$ for some $A + B + C = 0$.

Therefore, any element of $N(\widetilde{\mathrm{ch}})^{\mathbf{T}}$ can be *uniquely* represented by stacky S-families $\{\hat{F}_i\}_{i=1,2,3}$ of a μ-stable rank 2 toric vector bundle as in Proposition 7.5 satisfying

$$\begin{aligned}&\widetilde{\mathrm{ch}}(\mathcal{F}) = \widetilde{\mathrm{ch}},\ A_2 = A_3 = 0,\ p_1 = (1:0),\ p_2 = (0:1),\ p_3 = (1:1),\\ &0 < \Delta_1 < \Delta_2 + \Delta_3,\ 0 < \Delta_2 < \Delta_1 + \Delta_3,\ 0 < \Delta_3 < \Delta_1 + \Delta_2.\end{aligned}$$

Here we have used Proposition 7.7 to characterise μ-stability. The result now follows from Lemma 7.8. $\square$

The generating function of the previous theorem is the most refined version possible. Easier formulae can be obtained by specializing. One such specialization is grouping together moduli spaces with the same value of the constant term of the modified Hilbert polynomial $P_{\mathcal{O}_{\mathbf{P}}(1),\mathcal{E}}(-,t)$. In this context, it is more natural to fix $\alpha = \widetilde{\mathrm{ch}}^2(-\otimes\mathcal{E}) \in A^0(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ and $\beta = \widetilde{\mathrm{ch}}^1(-\otimes\mathcal{E}) \in A^1(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ as our topological invariants. In other words, we are interested in the Euler characteristic of the moduli space of μ-stable rank 2 vector bundles $\mathcal{F}$ on $\mathbb{P}$ with $\widetilde{\mathrm{ch}}^2(\mathcal{F}\otimes\mathcal{E}) = \alpha$, $\widetilde{\mathrm{ch}}^1(\mathcal{F}\otimes\mathcal{E}) = \beta$ and arbitrarily fixed $P_{\mathcal{O}_{\mathbf{P}}(1),\mathcal{E}}(\mathcal{F},0) \in \mathbb{Z}$. We denote by q the formal variable corresponding to $P_{\mathcal{O}_{\mathbf{P}}(1),\mathcal{E}}(\mathcal{F},0)$ and denote the corresponding generating function by $\bar{\mathsf{H}}^{vb}_{\alpha,\beta}(q)$.

Let $\mathcal{F}$ be a μ-stable rank 2 toric vector bundle on $\mathbb{P}$ described, as above, by the data $A_1 = A_2 = 0$, $A_3 = A \in \mathbb{Z}$, $\Delta_1, \Delta_2, \Delta_3 \in \mathbb{Z}_{>0}$ satisfying $b \mid \Delta_1$, $c \mid \Delta_2$, $a \mid \Delta_3$. Define $\phi_E(x) := x + 2x^2 + 3x^3 + \cdots + (E-1)x^{E-1}$. Then

$$\begin{aligned}&\widetilde{\mathrm{ch}}^2(\mathcal{F}\otimes\mathcal{E})_0 = 2E,\\ &\widetilde{\mathrm{ch}}^2(\mathcal{F}\otimes\mathcal{E})_f = 0,\ \forall f \in D\setminus 0,\\ &\widetilde{\mathrm{ch}}^1(\mathcal{F}\otimes\mathcal{E})_0 = -(2A + \Delta_1 + \Delta_2 + \Delta_3 - (E-1))E,\\ &\widetilde{\mathrm{ch}}^1(\mathcal{F}\otimes\mathcal{E})_f = 2\phi_E(e^{2\pi\sqrt{-1}f})e^{-2\pi\sqrt{-1}fA},\ \forall f \in D\setminus 0,\\ &\widetilde{\mathrm{ch}}^1(\mathcal{F}\otimes\mathcal{E})_f = 0,\ \forall f \in D_{12}, D_{13}, D_{23}.\end{aligned}$$

Here the vanishing in the second and last line are due to the presence of the generating sheaf $\mathcal{E}$. From these equations, we see that the classes α, β need to satisfy corresponding vanishing and divisorial properties or otherwise $\bar{\mathsf{H}}^{vb}_{\alpha,\beta}(q)$ is simply zero. In particular. β_0, $0 \in D$ has to be divisible by E. If we define $c_1 := \frac{\beta_0}{E} - (E-1)E \in \mathbb{Z}$ and $\lambda \in \{0,\ldots,d-1\}$, then fixing α and β is equivalent to fixing c_1, λ and the previous equations are equivalent to

$$-2A - \Delta_1 - \Delta_2 - \Delta_3 = c_1 \text{ and } A \equiv \lambda \bmod d. \tag{10}$$

These equations should be thought of as fixing "the first Chern class" and the $\boldsymbol{\mu}_d$-eigenvalue of the rank 2 vector bundles we are considering.

We now compute $P_{\mathcal{E}}(\mathcal{F},0)$. First some necessary notation. For any integers m and $n>0$, we denote by $[m]_n$ the unique value $k \in \{0,\ldots,n-1\}$ for which $m \equiv k \bmod n$. Also, for integers m_1, m_2, m_3 and $n>0$ define

$$\psi_E(m_1,m_2,m_3,n) := \sum_{\substack{k=1 \\ \frac{n}{\gcd(m_1,n)} \nmid k}}^{n-1} \frac{1+e^{-\frac{2\pi\sqrt{-1}km_2}{n}}}{1-e^{\frac{2\pi\sqrt{-1}km_1}{n}}} e^{-\frac{2\pi\sqrt{-1}km_3}{n}} \phi_E(e^{\frac{2\pi\sqrt{-1}k}{n}}).$$

Lemma 7.8 and the expression for the Todd classes in Section 4.2 imply that $P_{\mathcal{E}}(\mathcal{F},0) \in \mathbb{Z}$ is given by

$$\begin{aligned}
&\frac{E}{abc}\Big\{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1+\frac{1}{2}(a+b+c)c_1\\
&+\frac{1}{6}(a^2+b^2+c^2)+\frac{1}{2}(ab+bc+ca)+(c_1+a+b+c+E-d)[A]_d\\
&+\frac{1}{2}(c_1+a+b+c)(E-d)+[A]_d^2+\frac{1}{3}E^2-\frac{1}{2}Ed+\frac{1}{6}d^2\Big\}\\
&+\frac{1}{ab}\psi_E(c,\Delta_2,A,d_{12})+\frac{1}{ac}\psi_E(b,\Delta_1,A,d_{13})+\frac{1}{bc}\psi_E(a,\Delta_3,A,d_{23}).
\end{aligned}$$

For fixed c_1, λ, the above expression is a function the Δ_i only. Recall that A is determined by Equation (10). We denote this function by $P(\Delta_1,\Delta_2,\Delta_3)$. The generating function $\bar{\mathsf{H}}^{vb}_{c_1,\lambda}(q)$ is given by the following explicit formula

$$\sum_{\substack{\Delta_1,\Delta_2,\Delta_3 \in \mathbb{Z}_{>0} \\ b \mid \Delta_1,\, c \mid \Delta_2,\, a \mid \Delta_3 \\ 2 \mid c_1+\Delta_1+\Delta_2+\Delta_3 \\ -\frac{1}{2}(c_1+\Delta_1+\Delta_2+\Delta_3) \equiv \lambda \bmod d \\ \Delta_1 < \Delta_2+\Delta_3 \\ \Delta_2 < \Delta_1+\Delta_3 \\ \Delta_3 < \Delta_1+\Delta_2}} q^{P(\Delta_1,\Delta_2,\Delta_3)}.$$

In the following examples, $0 < \Delta_i < \Delta_j + \Delta_k$ is short hand for $0 < \Delta_1 < \Delta_2 + \Delta_3$, $0 < \Delta_2 < \Delta_1 + \Delta_3$ and $0 < \Delta_3 < \Delta_1 + \Delta_2$. In examples 1–3, it is easy to see that if $\gcd(2,c_1)=1$, then there are no strictly μ-semistables around. In example 4, a slightly more complicated numerical criterion is given.

Example 1. Let $\mathbb{P} = \mathbb{P}^2$, then $I\mathbb{P} = \mathbb{P}$. We take $\mathcal{E} = \mathcal{O}_{\mathbb{P}}$. Fixing $\alpha = 2$ and $\beta = c_1 = (\widetilde{\mathrm{ch}}^1)_0$, $0 \in D$, the generating function $\mathsf{H}^{vb}_{\alpha,\beta}$ is given by

$$\sum_{\substack{0 < \Delta_i < \Delta_j + \Delta_k \\ 2 \mid c_1 + \Delta_1 + \Delta_2 + \Delta_3}} p^{\frac{1}{4}c_1^2 + \frac{1}{4}\Delta_1^2 + \frac{1}{4}\Delta_2^2 + \frac{1}{4}\Delta_3^2 - \frac{1}{2}\Delta_1\Delta_2 - \frac{1}{2}\Delta_2\Delta_3 - \frac{1}{2}\Delta_3\Delta_1}.$$

Replacing p by q, this is also equal to the generating function $\bar{\mathsf{H}}^{vb}_{c_1,0}(q)$ (up to an additional constant $\frac{3}{2}c_1 + 2$ in the power of q making it integer-valued). This is essentially the generating function computed in [Koo2, Sect. 4.2.1]. It was first computed using localization methods by Klyachko [Kly2], who expresses it in terms of Hurwitz class numbers.

Example 2. Let $\mathbb{P} = \mathbb{P}(1,1,2)$, then $I\mathbb{P} = \mathbb{P} \sqcup \mathbb{P}(2)$. We take $\mathcal{E} = \mathcal{O}_{\mathbb{P}} \oplus \mathcal{O}_{\mathbb{P}}(1)$. The codegree 0 Chern character over the twisted sector $\mathbb{P}(2)$ is (Lemma 7.8)

$$(\widetilde{\mathrm{ch}}^0)_{1/2} = (-1)^A((-1)^{\Delta_1} + (-1)^{\Delta_3}).$$

Recall that $2 \mid \Delta_2$. This part of the Chern character can only take values $-2, 0, 2$. We denote the variable keeping track of $(\widetilde{\mathrm{ch}}^0)_0$, $0 \in D$ by p and the variable keeping track of $(\widetilde{\mathrm{ch}}^0)_{1/2}$, $1/2 \in D_3$ by r. For any subset $X \subset \mathbb{Z}^3$, we denote by $\mathbb{1}_X$ the characteristic function taking value 1 on X and 0 elsewhere. Fixing $\alpha = 2$ and $\beta = c_1 = (\widetilde{\mathrm{ch}}^1)_0$, $0 \in D$, the generating function $\mathsf{H}^{vb}_{\alpha,\beta}$ is

$$\sum_{\substack{0 < \Delta_i < \Delta_j + \Delta_k \\ 2 \mid c_1 + \Delta_1 + \Delta_2 + \Delta_3,\, 2 \mid \Delta_2}} p^{\frac{1}{4}c_1^2 + \frac{1}{4}\Delta_1^2 + \frac{1}{4}\Delta_2^2 + \frac{1}{4}\Delta_3^2 - \frac{1}{2}\Delta_1\Delta_2 - \frac{1}{2}\Delta_2\Delta_3 - \frac{1}{2}\Delta_3\Delta_1},$$

multiplied by

$$\begin{aligned}
&\left(\mathbb{1}_{\left\{\substack{4 \nmid c_1 + \Delta_1 + \Delta_2 + \Delta_3 \\ 2 \mid \Delta_1, 2 \mid \Delta_3}\right\}} + \mathbb{1}_{\left\{\substack{4 \mid c_1 + \Delta_1 + \Delta_2 + \Delta_3 \\ 2 \nmid \Delta_1, 2 \nmid \Delta_3}\right\}}\right) r^{-2} \\
&+ \left(\mathbb{1}_{\left\{2 \mid \Delta_1, 2 \nmid \Delta_3\right\}} + \mathbb{1}_{\left\{2 \nmid \Delta_1, 2 \mid \Delta_3\right\}}\right) r^0 \\
&+ \left(\mathbb{1}_{\left\{\substack{4 \mid c_1 + \Delta_1 + \Delta_2 + \Delta_3 \\ 2 \mid \Delta_1, 2 \mid \Delta_3}\right\}} + \mathbb{1}_{\left\{\substack{4 \nmid c_1 + \Delta_1 + \Delta_2 + \Delta_3 \\ 2 \nmid \Delta_1, 2 \nmid \Delta_3}\right\}}\right) r^2.
\end{aligned}$$

The specialized generating function $\bar{\mathsf{H}}^{vb}_{c_1,0}(q)$ is given by

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 2\,\mid\, c_1+\Delta_1+\Delta_2+\Delta_3,\, 2\,\mid\,\Delta_2}} q^{\frac{1}{4}c_1^2+\frac{5}{2}c_1+6+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1}.$$

Up to a constant depending on c_1 in the power of q, this has the same shape as in example 1 *except* for the additional division requirement $2 \mid \Delta_2$.

Example 3. Let $\mathbb{P}=\mathbb{P}(1,2,2)$, then $I\mathbb{P}=\mathbb{P}\sqcup\mathbb{P}(2,2)$. We take $\mathcal{E}=\mathcal{O}_{\mathbb{P}}\oplus\mathcal{O}_{\mathbb{P}}(1)$. This time, the part of $A^*(I\mathbb{P})_{\boldsymbol{\mu}_\infty}$ coming from the twisted sector $\mathbb{P}(2,2)$ has a codegree 1 and 0 part. As before, we fix $\alpha=2$, $\beta_0=c_1=(\widetilde{\mathrm{ch}}^{1})_0$, $0\in D$, but this time we also fix $\beta_{1/2}=(\widetilde{\mathrm{ch}}^{1})_{1/2}$, $1/2\in D_{23}$. The formula for $\beta_{1/2}$ is

$$\beta_{1/2}=(-1)^A(1+(-1)^{\Delta_3}),$$

so it can take values $-2,0,2$. Recall that $2\mid\Delta_1$, $2\mid\Delta_2$. Each case gives a different generating function. We denote the variable keeping track of $(\widetilde{\mathrm{ch}}^{0})_0$, $0\in D$ by p and the variable keeping track of $(\widetilde{\mathrm{ch}}^{0})_{1/2}$, $1/2\in D_{23}$ by q.

(I) Fix $\beta_{1/2}=-2$. This is equivalent to requiring $2\mid\Delta_3$ and $4\nmid c_1+\Delta_1+\Delta_2+\Delta_3$. In this case, $(\widetilde{\mathrm{ch}}^{0})_{1/2}$, $1/2\in D_{23}$ is simply given by

$$\begin{aligned}(\widetilde{\mathrm{ch}}^{0})_{1/2}&=-(-1)^A\left((1+(-1)^{\Delta_3})A+\Delta_1+\Delta_2+(-1)^{\Delta_3}\Delta_3\right)\\&=2A+\Delta_1+\Delta_2+\Delta_3=-c_1,\end{aligned}$$

so the generating function $\mathsf{H}^{vb}_{\alpha,\beta}$ is given by

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 2\,\mid\, c_1+\Delta_1+\Delta_2+\Delta_3 \\ 4\,\nmid\, c_1+\Delta_1+\Delta_2+\Delta_3 \\ 2\,\mid\,\Delta_1,2\,\mid\,\Delta_2,2\,\mid\,\Delta_3}} p^{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1}q^{-c_1}.$$

(II) Fix $\beta_{1/2}=0$. This is equivalent to requiring $2\nmid\Delta_3$. In this case, the formula for $(\widetilde{\mathrm{ch}}^{0})_{1/2}$, $1/2\in D_{23}$ is

$$\begin{aligned}(\widetilde{\mathrm{ch}}^{0})_{1/2}&=-(-1)^A\left((1+(-1)^{\Delta_3})A+\Delta_1+\Delta_2+(-1)^{\Delta_3}\Delta_3\right)\\&=(-1)^{\frac{1}{2}(c_1+\Delta_1+\Delta_2+\Delta_3)}(-\Delta_1-\Delta_2+\Delta_3).\end{aligned}$$

The generating function $\mathsf{H}^{vb}_{\alpha,\beta}$ is given by

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 2 \mid c_1+\Delta_1+\Delta_2+\Delta_3 \\ 2 \mid \Delta_1, 2 \mid \Delta_2, 2 \nmid \Delta_3}} p^{\substack{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2 \\ -\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1}} q^{(-1)^{\frac{1}{2}(c_1+\Delta_1+\Delta_2+\Delta_3)}(-\Delta_1-\Delta_2+\Delta_3)},$$

where we wrote the power of p over two lines.

(III) Fix $\beta_{1/2}=2$. Similar to case I, we obtain

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 4 \mid c_1+\Delta_1+\Delta_2+\Delta_3 \\ 2 \mid \Delta_1, 2 \mid \Delta_2, 2 \mid \Delta_3}} p^{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1} q^{c_1}.$$

The specialized generating function $\bar{\mathsf{H}}^{vb}_{c_1,0}(q)$ is given by

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 2 \mid c_1+\Delta_1+\Delta_2+\Delta_3, 2 \mid \Delta_1, 2 \mid \Delta_2}} q^{\substack{\frac{1}{8}c_1^2+\frac{3}{2}c_1+\frac{17}{4}+\frac{1}{8}\Delta_1^2+\frac{1}{8}\Delta_2^2+\frac{1}{8}\Delta_3^2-\frac{1}{4}\Delta_1\Delta_2-\frac{1}{4}\Delta_2\Delta_3 \\ -\frac{1}{4}\Delta_3\Delta_1-\frac{1}{8}(-1)^{\frac{1}{2}(c_1+\Delta_1+\Delta_2+\Delta_3)}(1+(-1)^{\Delta_3})}}.$$

Example 4. Let $\mathbb{P}=\mathbb{P}(2,2,2)$, then $I\mathbb{P}=\mathbb{P}\sqcup\mathbb{P}$. We take $\mathcal{E}=\mathcal{O}_{\mathbb{P}}\oplus\mathcal{O}_{\mathbb{P}}(1)$. This time, we fix $(\widetilde{\mathrm{ch}}^{2})_f$ and $(\widetilde{\mathrm{ch}}^{1})_f$ for $f\in D=\{0,1/2\}$. We fix $\alpha_0=2$ and fixing $\alpha_{1/2}=(\widetilde{\mathrm{ch}}^{2})_{1/2}=\pm 2$ is equivalent to fixing the parity of A. Clearly, the generating function is zero unless $\beta_0=c_1=(\widetilde{\mathrm{ch}}^{1})_0=(-1)^A(\widetilde{\mathrm{ch}}^{1})_{1/2}=(-1)^A\beta_{1/2}$, so we assume this is the case. We use the formal variable p_0 to keep track of $(\widetilde{\mathrm{ch}}^{0})_0$ and p_1 to keep track of $(\widetilde{\mathrm{ch}}^{0})_{1/2}$.

(I) Fix $\alpha_{1/2}=-2$. Then $\mathsf{H}^{vb}_{\alpha,\beta}$ is

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 2 \mid c_1+\Delta_1+\Delta_2+\Delta_3 \\ 4 \nmid c_1+\Delta_1+\Delta_2+\Delta_3 \\ 2 \mid \Delta_1,\ 2 \mid \Delta_2,\ 2 \mid \Delta_3}} (p_0p_1^{-1})^{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1}.$$

(II) Fix $\alpha_{1/2}=2$. Then $\mathsf{H}^{vb}_{\alpha,\beta}$ is

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ 4 \mid c_1+\Delta_1+\Delta_2+\Delta_3 \\ 2 \mid \Delta_1,\ 2 \mid \Delta_2,\ 2 \mid \Delta_3}} (p_0p_1)^{\frac{1}{4}c_1^2+\frac{1}{4}\Delta_1^2+\frac{1}{4}\Delta_2^2+\frac{1}{4}\Delta_3^2-\frac{1}{2}\Delta_1\Delta_2-\frac{1}{2}\Delta_2\Delta_3-\frac{1}{2}\Delta_3\Delta_1}.$$

Similarly, for the specialized generating function $\bar{\mathsf{H}}^{vb}_{c_1,\lambda}(q)$, we fix $c_1 \in \mathbb{Z}$ and a $\boldsymbol{\mu}_2$-eigenvalue $\lambda \in \{0,1\}$. Note that the generating function is zero unless c_1 is even, so assume this is the case. We get the following expression for $\bar{\mathsf{H}}^{vb}_{c_1,\lambda}(q)$

$$\sum_{\substack{0<\Delta_i<\Delta_j+\Delta_k \\ -\frac{1}{2}(c_1+\Delta_1+\Delta_2+\Delta_3)\equiv\lambda \bmod 2 \\ 2\mid\Delta_1,2\mid\Delta_2,2\mid\Delta_3}} q^{\frac{1}{16}c_1^2+\frac{7}{8}c_1+\frac{23}{8}-\frac{1}{8}(-1)^{\lambda}(c_1+7)+\frac{1}{16}\Delta_1^2+\frac{1}{16}\Delta_2^2+\frac{1}{16}\Delta_3^2-\frac{1}{8}\Delta_1\Delta_2-\frac{1}{8}\Delta_2\Delta_3-\frac{1}{8}\Delta_3\Delta_1}.$$

Note that the choices $(\frac{1}{2}c_1,\lambda)=(\mathrm{even},0),(\mathrm{odd},1)$ give the $c_1=0$ generating function of example 1 and the choices $(\frac{1}{2}c_1,\lambda)=(\mathrm{odd},0),(\mathrm{even},1)$ give the $c_1=1$ generating function of example 1 (up to a constant in the power of q). It is not hard to see that for the choices $(\frac{1}{2}c_1,\lambda)=(\mathrm{odd},0),(\mathrm{even},1)$, there are no strictly μ-semistables around.

References

[BCS] L. A. Borisov, L. Chen and G. G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*, J. Amer. Math. Soc. 18 193–215 (2005).

[BH] L. A. Borisov and R. P. Horja, *On the K-theory of smooth toric DM stacks*, Snowbird lectures on string geometry 21–42, Contemp. Math. 401, Amer. Math. Soc., Providence RI (2006).

[DS] R. Dijkgraaf and P. Sułkowski, *Instantons on ALE spaces and orbifold partitions*, JHEP 0803 (2008) 013.

[FMN] B. Fantechi, E. Mann and F. Nironi, *Smooth toric DM stacks*, Crelle 648 201–244 (2010).

[Ful] W. Fulton, *Introduction to toric varieties*, Princeton University Press (1993).

[Got] L. Göttsche, *Theta functions and Hodge numbers of moduli spaces of sheaves on rational surfaces*, Comm. Math. Phys. 206 105–136 (1999).

[Gut] A. Gutwirth, *The action of an algebraic torus on the affine plane*, Trans. AMS 105 407–414 (1962).

[Har] R. Harthorne, *Algebraic geometry*, Springer-Verlag (1977).

[Kan] T. Kaneyama, *On equivariant vector bundles on an almost homogeneous variety*, Nagoya Math. J. 57 65–86 (1975).

[Kly1] A. A. Klyachko, *Equivariant bundles on toral varieties*, Math. USSR Izvestiya 35 337–375 (1990).

[Kly2] A. A. Klyachko, *Vector bundles and torsion free sheaves on the projective plane*, preprint Max Planck Institut für Mathematik (1991).

[Koo1] M. Kool, *Fixed point loci of moduli spaces of sheaves on toric varieties*, Adv. Math. 227 1700–1755 (2011).

[Koo2] M. Kool, *Euler characteristics of moduli spaces of torsion free sheaves on toric surfaces*, arXiv:0906.3393.

[KR1] M. Koras and P. Russell, *Contractible threefolds and $\mathbb{C}^*$-actions on $\mathbb{C}^3$*, J. Alg. Geom. 6 671–695 (1997).

[KR2] M. Koras and P. Russell, *$\mathbb{C}^*$-actions on $\mathbb{C}^3$: the smooth locus of the quotient is not of hyperbolic type*, J. Alg. Geom. 8 603–694 (1999).

[Nir] F. Nirono, *Moduli spaces of semistable sheaves on projective Deligne-Mumford stacks*, arXiv:0811.1949.

[OS] M. Olsson and J. Starr, *Quot functors for Deligne-Mumford stacks*, Comm. Alg. 31 2069–4096 (2003). Special issue in honour of Steven L. Kleiman.

[Pay] S. Payne, *Moduli of toric vector bundles*, Compos. Math. 144 1199–1213 (2008).

[Per1] M. Perling, *Graded rings and equivariant sheaves on toric varieties*, Math. Nachr. 263–264 181–197 (2004).

[Per2] M. Perling, *Moduli for equivariant vector bundles of rank two on smooth toric surfaces*, Math. Nachr. 265 87–99 (2004).

[Per3] M. Perling, *Resolutions and moduli for equivariant sheaves over toric varieties*, Ph.D. dissertation University of Kaiserslautern (2003).

[Rom] M. Romagny, *Group actions on stacks and applications*, Michigan Math. J. 53 209–236 (2005).

[RT] M. Rossi and L. Terracini, *Weighted projective spaces from the toric point of view with computational applications*, arXiv:1112.1677.

[Toe] B. Toën, *Théorèmes de Riemann-Roch pour les champs de Deligne-Mumford*, K-theory 18 33–76 (1999).

[Vis] A. Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*, Invent. Math. 97 613–670 (1989).

`amingh@math.umd.edu`
Department of Mathematics
Mathematics Building
University of Maryland
College Park, MD 20742-4015
USA

`y.jiang@imperial.ac.uk`
`m.kool@imperial.ac.uk`
Department of Mathematics
Imperial College London
South Kensington Campus
London SW7 2AZ
United Kingdom